## **New devices**

- Lee TS, Hines GL, Feuerman M. Significant correlation between cerebral oximetry and carotid stump pressure during carotid endarterectomy. *Ann Vasc Surg* 2008:22:58—62.
- Nemoto EM, Yonas H, Kassam A. Clinical experience with cerebral oximetry in stroke and cardiac arrest. Crit Care Med 2000;28:1052—4.
- Seifert H, Blass G, Leetz HK, et al. The radiation exposure of the patient from stablexenon computed tomography. Br J Radiol 1995;68:301—5.
- Latchaw RE, Yonas H, Pentheny SL, et al. Adverse reactions to xenon-enhanced CT cerebral blood flow determination. Radiology 1987;163:251—4.
- Hillman J, Sturnegk P, Yonas H, et al. Bedside monitoring of CBF with xenon-CT and a mobile scanner: a novel method in neurointensive care. Br J Neurosurg 2005;19:395—401
- DeWitt DS, Fatouros PP, Wist AO, et al. Stable xenon versus radiolabeled microsphere cerebral blood flow measurements in baboons. Stroke 1989:20:1716—23
- Fainardi E, Tagliaferri MF, Compagnone C, et al. Regional cerebral blood flow levels as measured by xenon-CT in vascular territorial low-density areas after subarachnoid hemorrhage are not always ischemic. Neuroradiology 2006;48:685—90.
- Yonas H, Pindzola RP, Johnson DW. Xenon/computed tomography cerebral blood flow and its use in clinical management. Neurosurg Clin N Am 1996;7:605—16
- Yonas H, Pindzola RR, Meltzer CC, et al. Qualitative versus quantitative assessment of cerebrovascular reserves. Neurosurgery 1998;42:1005—10.
- Yonas H, Good WF, Gur D, et al. Mapping cerebral blood flow by xenon-enhanced computed tomography: clinical experience. Radiology 1984;152:435–42.
- Firlik AD, Yonas H. The utility of XeCT cerebral blood flow in the management of acute stroke. Keio J Med 2000;49(Suppl 1):A129—30.

- Yonas H, Sekhar L, Johnson DW, et al. Determination of irreversible ischemia by xenon-enhanced computed tomographic monitoring of cerebral blood flow in patients with symptomatic vasospasm. Neurosurgery 1989;24:368—72.
- Inoue Y, Shiozaki T, Tasaki O, et al. Changes in cerebral blood flow from the acute to the chronic phase of severe head injury. J Neurotrauma 2005;22:1411–18.
- Firlik AD, Yonas H, Kaufmann AM, et al. Relationship between cerebral blood flow and the development of swelling and life-threatening herniation in acute ischemic stroke. J Neurosurg 1998;89:243—9.
- Gupta R, Crago EA, Gallek M, et al. Reduced ipsilateral hemispheric cerebral blood flow at admission is predictive of vasospasm with infarction after aneurysmal subarachnoid hemorrhage. Neurocrit Care 2008;9:27—30.
- Wintermark M, Thiran JP, Maeder P, et al. Simultaneous measurement of regional cerebral blood flow by perfusion CT and stable xenon CT: a validation study. AJNR Am J Neuroradiol 2001;22:905—14.
- Hashimoto K, Murakami T, Dono K, et al. Quantitative tissue blood flow measurement of the liver parenchyma: comparison between xenon CT and perfusion CT. Dia Dis Sci 2007;52:943—9.
- Thavasothy M, Broadhead M, Elwell C, et al. A comparison of cerebral oxygenation as measured by the NIRO 300 and the INVOS 5100 near-infrared spectrophotometers. Anaesthesia 2002;57:999—1006.
- Torella F, Cowley R, Thorniley MS, et al. Monitoring blood loss with near infrared spectroscopy. Comp Biochem Physiol A Mol Integr Physiol 2002;132:199—203.
- Torella F, Cowley RD, Thorniley MS, et al. Regional tissue oxygenation during hemorrhage: can near infrared spectroscopy be used to monitor blood loss? Shock 2002:18:440—4.
- Torella F, Haynes SL, McCollum CN. Cerebral and peripheral near-infrared spectroscopy: an alternative transfusion trigger? Vox Sang 2002;83:254-7.

## Correction

Jensen ME, McGraw JK, Cardella JF, et al. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the American Society of Interventional and Therapeutic Neuroradiology, Society of Interventional Radiology, American Association of Neurological Surgeons/Congress of Neurological Surgeons, and American Society of Spine Radiology. J NeuroInterv Surg 2009;1:181–5.

The doi number used for this article in December 2009 Volume 1 Issue 2 should not have been used. The correct doi number should be: doi:10.1136/j.jvir.2009.04.022.

J NeuroIntervent Surg 2011;3:398. doi:10.1136/j.jvir.2009.04.022corr1