



Editor's choice
Scan to access more
free content

ORIGINAL RESEARCH

Predictors of poor outcome despite recanalization: a multiple regression analysis of the NASA registry

Italo Linfante,^{1,2} Amy K Starosciak,^{2,3} Gail R Walker,³ Guilherme Dabus,^{1,2} Alicia C Castonguay,⁴ Rishi Gupta,⁵ Chung-Huan J Sun,⁵ Coleman Martin,⁶ William E Holloway,⁶ Nils Mueller-Kronast,⁷ Joey D English,⁸ Tim W Malisch,⁹ Franklin A Marden,⁹ Hormozd Bozorgchami,¹⁰ Andrew Xavier,¹¹ Ansaar T Rai,¹² Michael T Froehler,¹³ Aamir Badruddin,¹⁴ Thanh N Nguyen,¹⁵ M Asif Taqi,⁶ Michael G Abraham,¹⁷ Vallabh Janardhan,¹⁸ Hashem Shaltoni,¹⁹ Roberta Novakovic,²⁰ Albert J Yoo,²¹ Alex Abou-Chebl,²² Peng R Chen,²³ Gavin W Britz,²⁴ Ritesh Kaushal,²⁵ Ashish Nanda,²⁶ Mohammad A Issa,⁴ Raul G Nogueira,⁵ Osama O Zaidat⁴

For numbered affiliations see end of article.

Correspondence to

Dr Italo Linfante, Miami Cardiac and Vascular Institute and Neuroscience Center, Baptist Hospital, Miami, FL, USA; linfante.italo@gmail.com

Received 24 October 2014
Revised 9 December 2014
Accepted 12 December 2014
Published Online First
6 January 2015

ABSTRACT

Background Mechanical thrombectomy with stent-retrievers results in higher recanalization rates compared with previous devices. Despite successful recanalization rates (Thrombolysis in Cerebral Infarction (TICI) score $\geq 2b$) of 70–83%, good outcomes by 90-day modified Rankin Scale (mRS) score ≤ 2 are achieved in only 40–55% of patients. We evaluated predictors of poor outcomes (mRS > 2) despite successful recanalization (TICI $\geq 2b$) in the North American Solitaire Stent Retriever Acute Stroke (NASA) registry.

Methods Logistic regression was used to evaluate baseline characteristics and recanalization outcomes for association with 90-day mRS score of 0–2 (good outcome) vs 3–6 (poor outcome). Univariate tests were carried out for all factors. A multivariable model was developed based on backwards selection from the factors with at least marginal significance ($p \leq 0.10$) on univariate analysis with the retention criterion set at $p \leq 0.05$. The model was refit to minimize the number of cases excluded because of missing covariate values; the c-statistic was a measure of predictive power.

Results Of 354 patients, 256 (72.3%) were recanalized successfully. Based on 234 recanalized patients evaluated for 90-day mRS score, 116 (49.6%) had poor outcomes. Univariate analysis identified an increased risk of poor outcome for age ≥ 80 years, occlusion site of internal carotid artery (ICA)/basilar artery, National Institute of Health Stroke Scale (NIHSS) score ≥ 18 , history of diabetes mellitus, TICI 2b, use of rescue therapy, not using a balloon-guided catheter or intravenous tissue plasminogen activator (IV t-PA), and > 30 min to recanalization ($p \leq 0.05$). In multivariable analysis, age ≥ 80 years, occlusion site ICA/basilar, initial NIHSS score ≥ 18 , diabetes, absence of IV t-PA, ≥ 3 passes, and use of rescue therapy were significant independent predictors of poor 90-day outcome in a model with good predictive power (c-index=0.80).

Conclusions Age, occlusion site, high NIHSS, diabetes, no IV t-PA, ≥ 3 passes, and use of rescue therapy are associated with poor 90-day outcome despite successful recanalization.

INTRODUCTION

Recanalization of the occluded artery is a powerful predictor of good outcome in acute ischemic stroke secondary to large artery occlusion.^{1–4} Mechanical thrombectomy with stent-retrievers results in higher recanalization rates and better outcomes than previous devices such as the Concentric Thrombus Retriever. However, despite rates of successful recanalization (Thrombolysis in Cerebral Infarction (TICI) score $\geq 2b$) up to 85%, good clinical outcomes assessed by modified Rankin Scale (mRS) ≤ 2 are achieved in only up to 55% of patients.^{2–7}

Some authors have identified factors that influence poor outcomes in patients with acute stroke treated with the Merci thrombectomy device. In particular, the multi MERCI trial identified absence of successful recanalization, age, high National Institute of Health Stroke Scale (NIHSS) score, and proximal vessel occlusion as predictors of mortality.^{1, 8} However, it is not clear which factors increase the risk of a poor clinical outcome despite recanalization.

The North American Solitaire Acute Stroke (NASA) registry is a multicenter, non-sponsored, physician-conducted, post-marketing registry on the use of the Solitaire FR device in 354 patients with acute large vessel ischemic stroke. Recanalization rates and clinical outcomes reported in NASA⁷ were comparable to the data from the randomized trials SWIFT⁵ and TREVO.⁶

The present study evaluated baseline characteristics and recanalization parameters for association with poor outcomes (mRS ≥ 3) in successfully recanalized (TICI $\geq 2b$) cases from the NASA registry of patients with acute stroke treated with the Solitaire FR device.

METHODS

Research participants

Study participants (de-identified data) were obtained from the NASA registry of patients treated with the Solitaire FR as the only device for restoration of blood flow. The NASA registry recruited



CrossMark

To cite: Linfante I, Starosciak AK, Walker GR, et al. *J NeuroIntervent Surg* 2016;**8**:224–229.

24 clinical sites within North America to submit retrospective demographic, clinical presentation, site-adjudicated angiographic, procedural, and clinical outcome data on consecutive patients with acute stroke treated within 8 h of symptom onset with the Solitaire FR device from March 2012 to February 2013. Details of the NASA participant population can be found in the original report.⁷

Data

In addition to patient demographics, registry data included information on revascularization and clinical outcomes. Thrombolysis in Myocardial Infarction (TIMI) and TIC1 scores were determined based on the final angiogram after the procedure was completed. Successful recanalization was defined as TIC1 $\geq 2b$. Clinical outcomes included whether or not patients developed a symptomatic intracranial hemorrhage upon 24 h CT follow-up; mRS score at discharge, 30 and 90 days post-treatment; NIHSS score at pretreatment, discharge, and 90 days post-treatment; and mortality. A good clinical outcome was defined as mRS ≤ 2 at 90 days.

Statistical analysis

Descriptive statistics were used to characterize study patients at baseline and recanalization outcomes. Counts and percentages are reported for categorical variables; continuous variables are summarized as mean (SD) or median (range). Baseline characteristics included demographics, comorbidities, site of stroke, and initial NIHSS score. Recanalization outcomes included post-procedure TIC1 score, number of passes, use of intravenous tissue plasminogen activator (IV t-PA), balloon-guided catheter (BGC), Penumbra or rescue therapy, time from symptom onset to treatment start, and time from treatment start to recanalization.

Logistic regression was used to evaluate baseline characteristics and recanalization results for association with 90-day outcomes characterized as good (mRS 0–2) or poor (mRS 3–6). First, univariate tests were carried out for all factors except time from onset to treatment initiation (see below). Age and time to recanalization were modeled categorically after determining that the linearity assumption (constant risk per unit change) did not hold. Category cut-off points for these variables were then derived from data quartiles. Quartile analysis was also used for initial testing of NIHSS.

Next, a multivariable model was developed based on backwards selection from the set of factors with at least marginal significance ($p \leq 0.10$) on univariate analysis. The retention criterion was set at $p \leq 0.05$. The resulting model was refit to minimize the number of cases excluded because of missing covariate values, and the c-statistic was used as a measure of predictive power. The sensitivity of model results to missing outcome data was also evaluated. This analysis was done using the most recent available mRS score (last observation carried forward (LOCF) method) and also under the worst case assumption of poor outcome for all cases with missing 90-day mRS scores.

Time from onset to treatment initiation was evaluated in a separate analysis restricted to the subset of anterior site cases treated within 8 h of symptom onset. The restriction was necessary because posterior site cases tended to have less urgent symptoms and greater tolerance of delayed treatment.

Statistical analysis was carried out using SAS software V.9.3 (SAS Institute, Cary, North Carolina, USA).

RESULTS

The NASA registry enrolled 354 consecutive patients with acute ischemic stroke treated with the Solitaire FR device.⁷ Of these,

256 (72.3%) were successfully recanalized (TIC1 2b–3) and 234 recanalized patients had 90-day mRS scores as required entry criteria for our analysis. Baseline characteristics and recanalization outcomes for study patients are summarized in table 1.

At 90-day follow-up, 118 (50.4%) patients were classified as having a good outcome (mRS 0–2) while 116 (49.6%) had a poor outcome (mRS 3–6). An analysis of factors associated with a poor outcome following recanalization is presented in table 2. Univariate tests identified an increased risk for each of the following: age ≥ 80 years (upper quartile of data), internal carotid artery (ICA)/basilar site, median initial NIHSS score ≥ 18 , diabetes mellitus (DM), final TIC1 2b, BGC not used, IV t-PA not used, time to recanalization (≥ 30 min vs < 30 min), and use of rescue therapy ($p < 0.05$). No association was found for sex, race, smoking, atrial fibrillation, coronary artery disease, hypertension,

Table 1 Baseline characteristics and recanalization outcomes for 234 patients

Baseline characteristics	Mean N	SD %
Mean age, years	66.9	14.7
Sex (1)		
Female	119	51.1
Male	114	48.9
Race (1)		
White	179	76.8
Black	39	16.7
Other	15	6.4
Smoking (3)	67	29.0
Atrial fibrillation	98	41.9
Diabetes mellitus	61	26.1
Hypertension	174	74.4
Hyperlipidemia	124	53.0
Coronary artery disease	76	32.5
Location of clot/occlusion		
M1	136	58.1
M2	23	9.8
ICA	48	20.5
Basilar	27	11.5
NIHSS, baseline (6)		
Mild: 0–5	10	4.4
Moderate: 6–19	127	55.7
Severe: 20–42	91	39.9
Recanalization outcomes		
Final TIC1 score		
2b	107	45.7
3	127	54.3
Rescue therapy, Solitaire failed	48	20.5
Distal embolization (2)	36	15.5
Balloon-guided catheter (10)	98	43.8
Solitaire+Penumbra (10)	32	14.3
Intravenous t-PA (1)	100	42.9
Intra-arterial t-PA	65	27.8
Passes (1)		
1–2	198	84.9
≥ 3	35	15.0
Median (range) time from onset to procedure start (8), min	277	75–1425
Median (range) time from procedure start to recanalization (25), min	45	8–412

Data shown are n (%) unless indicated otherwise.

ICA, internal carotid artery; NIHSS, National Institutes of Health Stroke Scale; t-PA, tissue plasminogen activator.

Table 2 Effect of baseline characteristics and recanalization outcomes on risk of poor 90-day outcome (mRS 3–6)

	N (%) with risk factor		Crude OR	95% CI	p Value	Adjusted OR	95% CI	p Value
	Good (mRS 0–2)	Poor (mRS 3–6)						
Baseline characteristics								
Age ≥80 vs <80 years	20/118 (17)	33/116 (28)	1.95	(1.04 to 3.65)	0.037	2.50	(1.21 to 5.19)	0.014
Male vs female	52/117 (44)	62/116 (53)	1.44	(0.86 to 2.40)	0.170	–	–	–
Black vs white	19/117 (16)	20/116 (17)	1.09	(0.54 to 2.18)	0.810	–	–	–
Other race vs white	7/117 (6)	8/116 (7)	1.18	(0.41 to 3.40)	0.757	–	–	–
Smoking	39/117 (33)	28/114 (25)	0.65	(0.37 to 1.16)	0.143	–	–	–
Atrial fibrillation	46/118 (39)	52/116 (45)	1.27	(0.76 to 2.14)	0.366	–	–	–
Diabetes mellitus	22/118 (19)	39/116 (34)	2.21	(1.21 to 4.04)	0.010	2.82	(1.37 to 5.83)	0.005
Hypertension	85/118 (72)	89/116 (77)	1.28	(0.71 to 2.31)	0.412	–	–	–
Hyperlipidemia	61/118 (52)	63/116 (54)	1.11	(0.66 to 1.86)	0.689	–	–	–
Coronary artery disease	34/118 (29)	42/116 (36)	1.40	(0.81 to 2.43)	0.228	–	–	–
ICA/basilar site vs M1/M2	29/118 (25)	46/116 (40)	2.02	(1.15 to 3.53)	0.014	2.18	(1.10 to 4.33)	0.026
NIHSS ≥18 vs ≤17	42/117 (36)	73/111 (66)	3.43	(1.99 to 5.91)	<0.001	4.51	(2.33 to 8.71)	<0.001
Recanalization outcomes								
Final TICI 2b vs 3	46/118 (39)	61/116 (53)	1.74	(1.03 to 2.92)	0.037	–	–	–
Rescue therapy, Solitaire failed	12/118 (10)	36/116 (31)	3.97	(1.94 to 8.12)	<0.001	4.94	(2.14 to 11.43)	<0.001
Distal embolization	20/116 (17)	16/116 (14)	0.77	(0.38 to 1.57)	0.469	–	–	–
Balloon-guided catheter not used	54/113 (48)	72/111 (65)	2.02	(1.18 to 3.45)	0.010	–	–	–
Solitaire+Penumbra	14/113 (12)	18/111 (16)	1.37	(0.64 to 2.91)	0.414	–	–	–
Intra-arterial t-PA not used	89/118 (75)	80/116 (69)	0.72	(0.41 to 1.29)	0.271	–	–	–
Intravenous t-PA not used	57/118 (48)	76/115 (66)	2.09	(1.23 to 3.54)	0.006	2.81	(1.47 to 5.36)	0.002
Passes ≥3 vs 1–2	13/117 (11)	22/116 (19)	1.87	(0.89 to 3.93)	0.097	2.62	(1.03 to 6.69)	0.043
Time to recanalization ≥30 min vs <30 min	69/105 (66)	86/104 (83)	2.49	(1.28 to 4.59)	0.006	–	–	–

p Value: Wald statistic. ORs estimate the risk of a poor 90-day outcome for patients with versus without the factor shown at left except where other categories are indicated. Crude ORs were estimated from univariate logistic regression; adjusted ORs from a multivariable model with the covariates shown (226 cases). ICA, internal carotid artery; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; TICI, Thrombolysis in Cerebral Infarction; t-PA, tissue plasminogen activator.

hyperlipidemia, distal embolization, use of Solitaire in association with the Penumbra, and intra-arterial (IA) t-PA.

In multivariable analysis, age ≥80 years, ICA/basilar site, initial NIHSS score ≥18, DM, absence of IV t-PA, ≥3 passes, and use of rescue therapy were significant independent predictors of poor 90-day outcome. The model, which used 226 cases, had a c-index of 0.80 (95% CI 0.74 to 0.86), indicating good predictive power.

Data were insufficient to consider the prognostic effect of blood pressure, embolization to new territory (only 4 cases), transfer and transfer type (45 missing values), general anesthesia (45 missing values), fluoroscopy time (36 missing values), or time from door to groin puncture (76 missing values). Development of symptomatic intracranial hemorrhage was not included in the regression analysis because it is not known until 24 h after procedure completion.

Sensitivity analysis with respect to missing 90-day mRS scores for 22 recanalized patients confirmed the prognostic effects shown in table 2, with some differences in statistical significance as follows. Under LOCF and worst case assumptions, site was at most marginally significant (OR 1.7, $p=0.110$ and OR 1.8, $p=0.068$, respectively). LOCF but not worst case assumptions also resulted in the loss of significance for number of passes (OR 1.7, $p=0.195$). These models included 246 and 247 cases, respectively.

Subset analysis of anterior cases (M1, M2, ICA) with respect to total time to recanalization was based on 154 cases for which treatment was started within 8 h of symptom onset. Total time was marginally significant ($p=0.055$), with an estimated 9% increased risk of poor outcome per 30 min increase in time from onset to recanalization.

DISCUSSION

A large body of literature shows that, in acute ischemic stroke, recanalization is a powerful predictor of good outcomes. Notably, a meta-analysis by Rha and Saver⁹ of 53 articles published between 1985 and 2002 reported data on 1774 patients evaluated for vessel recanalization. Based on 33 of the studies totaling 998 patients, their meta-analysis reported that good outcomes were more frequent in the recanalized group at 3 months (OR 4.43; 95% CI 3.32 to 5.91) and 3-month mortality was reduced (OR 0.24; 95% CI 0.16 to 0.35).⁹ Interestingly, mechanical thrombectomy in SWIFT, STAR and NASA yielded similar results—namely, a significant increase in the odds of a good outcome (OR 4.87; 95% CI 2.59 to 9.14) and a risk reduction of approximately 60% for mortality (OR 0.38; 95% CI 0.23 to 0.65). Thus, the benefit of recanalization is well established. It remains important to understand which factors influence poor outcomes despite adequate recanalization.

In our study of patients with acute stroke in the NASA registry with recanalization of the occluded artery, several factors emerged as predictors of poor outcomes (mRS ≥3). In particular, age ≥80 years, occlusion site other than M1/M2, NIHSS score ≥18, DM, absence of pretreatment with IV t-PA, ≥3 passes with the Solitaire device, and use of rescue therapy were identified as significant independent predictors of poor 90-day outcome. The strongest effects were NIHSS score and rescue therapy, both of which increased the risk of a poor outcome approximately fourfold whereas the increase estimated for each of the other factors was 2–3.

Age (>80 years) has previously been associated with poor outcome in patients treated with IV t-PA⁸ and intra-arterial therapy (IAT) in an analysis of the PROACT II data.^{10 11} It has

been hypothesized that age per se significantly impacts clinical outcomes after cerebral ischemia regardless of the presence of reperfusion. Very recently, Ribo *et al*¹² analyzed the relationship between outcomes and final infarct volume (FIV) in 214 patients stratified by age. Similar to the data reported by Yoo *et al*,¹³ Ribo *et al* also reported that a FIV of ≤ 49 mL predicts a mRS score of ≤ 2 in patients aged <70 years. Interestingly, the authors found that the FIV threshold for good outcomes decreased to 32.5 mL in patients aged 70–79 years and to 15.2 mL in patients aged ≥ 80 years. The oldest age group had the smallest proportion of patients at or below the age-adjusted FIV threshold (14–44% depending on the initial Alberta Stroke Program CT Score (ASPECTS) vs 29–65% in the two younger age strata). The data have to be considered in light of the fact that ischemic stroke in elderly patients without recanalization either by thrombectomy or after IV t-PA is associated with very high rates of death and severe disability.^{14–15} In particular, patients aged >80 years with acute ischemic stroke have higher risk-adjusted fatality, longer hospitalization, and are less likely to be discharged home than younger patients.¹⁵ These consistent findings may be explained by the very low tolerance of elderly brain to volumes of infarction that are otherwise tolerated in younger patients and the decreased plasticity of neuronal networks as age advances. Nevertheless, elderly patients should be considered for acute endovascular intervention because of their extremely poor outcomes without any treatment. However, one has to consider that, even after successful recanalization, their outcomes may not be comparable to younger patients.

A high NIHSS score on admission has previously been correlated with poor outcomes after IV and IAT by several studies.^{16–18} In our analysis of the NASA data the NIHSS score was predictive of poor outcomes despite recanalization, in line with previous reports.^{16–18} The results seem to indicate that the task of reversing a high neurological deficit at presentation is difficult to accomplish. These data also show the dilemma of considering patients with high NIHSS for IAT and their inclusion in clinical trials. In this regard, the SYNTHESIS trial concluded that endovascular treatment was not superior to IV t-PA in patients with acute stroke presenting with a median NIHSS score of 13.¹⁹ However, in the SYNTHESIS trial, mechanical thrombectomy was used only in 33.3% of patients and more than 30% of patients with acute stroke presented with a NIHSS score ≤ 10 .¹⁹ Interestingly, in IMS III, patients with a NIHSS score >20 had better outcomes with IAT than patients treated with IV t-PA alone.²⁰ This finding suggests that patients who present with severe neurological deficits (as indicated by a high NIHSS score) have even poorer outcomes with IV therapy or medical management than with thrombectomy.

Acute ICA or basilar artery occlusion are known to be associated with poor outcomes.²¹ In particular, distal ICA occlusion and/or tandem occlusions (ICA plus middle cerebral artery (MCA)) carry a worse prognosis than MCA occlusions.²² In addition, it has been reported that ICA occlusions have a poorer response to IV t-PA than MCA occlusions.^{23–24} In another series, Zaidat *et al*²⁵ examined patients with acute occlusion of the distal ICA treated with IAT/IV rt-PA. In their study the mortality rate was 50% despite complete recanalization (80% in the combined IV/IA thrombolysis group and 62% in the group treated with IAT alone). In particular, patients with ICA, MCA, and anterior cerebral artery (terminus T) occlusions were the least likely to respond to thrombolysis and only 38.9% of the patients showed moderate-to-good outcomes (90-day mRS ≤ 3). As a result, patients with proximal ICA occlusions or basilar

occlusions are excluded from most of the clinical trials on acute stroke intervention. The correlation may be the result of a high initial infarct volume with proximal arterial occlusion. It is conceivable that, in proximal arterial occlusion, there may be technical difficulties in recanalizing the occluded vessel which results in a longer time to achieve recanalization. In this study there was an increased median time to recanalization for sites other than M1/M2 (51 min vs 44 min); however, the difference was not statistically significant ($p=0.260$, Wilcoxon test).

The need for ≥ 3 passes of the Solitaire FR and the use of rescue therapy were both retained in our multivariable model, whereas longer time to achieve successful recanalization was eliminated despite its significance on univariate analysis. This is explained by the fact that both the number of passes and rescue therapy were associated with longer recanalization times. Median time to recanalization for cases that required ≥ 3 passes was 85 min compared with 41 min when 1–2 passes were sufficient. Similarly, median time to recanalization was 78 min versus 41 min depending on whether rescue therapy was needed. Both associations were significant ($p<0.001$, Wilcoxon test).

Dramatic time sensitivity of the cerebral tissue to ischemia in acute ischemic stroke secondary to large artery occlusion has been shown by Saver.²⁶ Time dependency to initiation of thrombolysis was already shown in a large meta-analysis of all IV t-PA trials²⁷ and by other authors.²⁸ The lack of benefit from acute stroke intervention in IMS III and SYNTHESIS trials may also be related to delayed time to recanalization and therefore futile reperfusion.^{16–19–20} Based on the IV t-PA data, there is a global effort to shorten as much as possible the time from symptom onset to intervention and door to needle. Similar to treatment with IV t-PA, Pereira *et al*²⁹ have shown better outcomes in patients treated with mechanical thrombectomy early compared with later. Thus, several authors have already highlighted the importance of shortening as much as possible the time from image to groin puncture and from puncture to recanalization. The result of this analysis shows that the time dependency of intervention is critical to avoid futile reperfusion, as indicated by poor outcomes despite recanalization. The data suggest that faster time to intervention is probably associated with improved clinical outcomes.

Very recently Shi *et al*³⁰ reported an analysis of factors that may predict poor outcomes despite recanalization. The authors reported pooled data from the multi MERCI, TREVO, and TREVO 2 trials. Although the aim of their study was similar to ours, there are differences in their data acquisition. In particular, they analyzed pooled data obtained from patients treated with different devices (ie, Concentric Retriever and Trevo) that have been shown to be different in effectiveness for recanalization. Shi *et al* also used different recanalization scales (ie, TIMI vs TICI). Recanalization in the NASA registry was obtained with the Solitaire FR and assessed by TICI score in all 354 patients. Nevertheless, similar to our analysis, they report that advanced age, high NIHSS score, and delay from symptom onset to recanalization are predictors of dependency despite recanalization. Interestingly, the authors report an increase of 11% in the odds of functional dependence (mRS >3) for every 30 min delay from symptom onset to endovascular intervention. In our analysis of the NASA data, total time to recanalization was marginally significant ($p=0.055$) on univariate analysis with an estimated 9% increased risk of poor outcome (mRS >2) per 30 min delay in endovascular intervention. This difference may be because they analyzed pooled data from patients treated with the Concentric Retriever and Trevo whereas we analyzed data from patients treated with Solitaire FR.

Limitations of the study

Our analysis has several limitations. In the NASA registry, assessment of reperfusion and clinical outcomes was obtained without a core laboratory or requirement of an independent adjudicator. Nevertheless, recanalization rates and outcomes in the NASA registry are strikingly similar to controlled trials on mechanical thrombectomy with stent-retrievers (ie, SWIFT and TREVO trials).^{5–7} It has been reported that ischemic core imaging,¹³ ASPECT score, and collateral circulation assessment may correlate with outcomes after thrombectomy.^{29–31} In the NASA registry the ASPECT score and collateral circulation assessment was not performed systematically so these variables, although of great interest, were not available on all patients and are not therefore included in the analysis. Finally, there are inherent limitations in a retrospective study based on registry data acquired according to pre-established specifications.

CONCLUSIONS

In acute ischemic stroke secondary to large artery occlusion, revascularization of the occluded artery is unquestionably a powerful predictor of good outcome. Nevertheless, with the current technology and logistics, unfavorable outcomes (mRS ≥ 3) occur in 49% of patients despite successful recanalization. In these patients, older age, presentation with severe neurological deficit, delays to reperfusion, diabetes, occlusion site, and not receiving IV t-PA are important factors that predict poor clinical outcomes. Reducing delays in reperfusion, considering all these factors, may be the best way to improve clinical outcomes and should be considered in clinical trials that evaluate IA thrombectomy.

Author affiliations

¹Miami Cardiac and Vascular Institute, Baptist Hospital, Miami, Florida, USA

²Neuroscience Center, Baptist Hospital, Miami, Florida, USA

³Center for Research and Grants, Baptist Health South Florida, Coral Gables, Florida, USA

⁴Medical College of Wisconsin/Froedtert Hospital, Milwaukee, Wisconsin, USA

⁵Emory University School of Medicine, Atlanta, Georgia, USA

⁶St. Luke's Kansas City, Kansas City, Missouri, USA

⁷Delray Medical Center, Delray Beach, Florida, USA

⁸California Pacific Medical Center, San Francisco, California, USA

⁹Alexian Brothers Medical Center, Elk Grove Village, Illinois, USA

¹⁰Oregon Health and Sciences, Portland, Oregon, USA

¹¹Wayne State University School of Medicine, Detroit, Michigan, USA

¹²West Virginia University Hospital, Morgantown, West Virginia, USA

¹³Vanderbilt University Medical Center, Nashville, Tennessee, USA

¹⁴Provena St. Joseph Medical Center, Joliet, Illinois, USA

¹⁵Boston Medical Center, Boston, Massachusetts, USA

¹⁶Desert Regional Medical Center, Palm Springs, California, USA

¹⁷University of Kansas Medical Center, Kansas City, Kansas, USA

¹⁸Texas Stroke Institute, Plano, Texas, USA

¹⁹Baylor College of Medicine, Houston, Texas, USA

²⁰UT Southwestern Medical Center, Dallas, Texas, USA

²¹Massachusetts General Hospital, Boston, Massachusetts, USA

²²Baptist Health Louisville, Louisville, Kentucky, USA

²³The University of Texas Medical School at Houston, Houston, Texas, USA

²⁴Methodist Neurological Institute, Houston, Texas, USA

²⁵St Louis University, St Louis, Missouri, USA

²⁶University of Missouri, Columbia, Missouri, USA

Contributors IL developed the idea for the manuscript and participated in the design, concept, and data gathering for the NASA registry. IL, AKS, and GRW participated in data analysis, writing, and meaningful editing on the manuscript. GD, ACC, RG, C-HJS, CM, WEH, NM-K, JDE, TWM, FAM, HB, AX, ATR, MTF, AB, TNN, MAT, MGA, VJ, HS, RN, AJY, AA-C, PRC, GWB, RK, AN, MAI, RGN, and OOO participated in the design, conception, and data gathering for the NASA registry, meaningful editing of the manuscript, and provided suggestions and feedback on the manuscript. All authors approved the final manuscript.

Competing interests None.

Ethics approval Ethics approval was provided by the Baptist Health South Florida Institutional Review Board and the ethics committee at each contributing institution.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement There are no additional unpublished data from this study. This was a new analysis of data from the North American Solitaire Stent Retriever Acute Stroke registry that was published previously in JNIS.⁷

REFERENCES

- Nogueira RG, Liebeskind DS, Sung G, *et al*. Predictors of good clinical outcomes, mortality, and successful revascularization in patients with acute ischemic stroke undergoing thrombectomy: pooled analysis of the Mechanical Embolus Removal in Cerebral Ischemia (MERCI) and Multi MERCI Trials. *Stroke* 2009;40:3777–83.
- Smith WS. Safety of mechanical thrombectomy and intravenous tissue plasminogen activator in acute ischemic stroke. Results of the multi Mechanical Embolus Removal in Cerebral Ischemia (MERCI) trial, part I. *AJNR Am J Neuroradiol* 2006;27:1177–82.
- Nogueira RG, Smith WS. Safety and efficacy of endovascular thrombectomy in patients with abnormal hemostasis: pooled analysis of the MERCI and multi MERCI trials. *Stroke* 2009;40:516–22.
- Smith WS, Sung G, Saver J, *et al*. Mechanical thrombectomy for acute ischemic stroke: final results of the Multi MERCI trial. *Stroke* 2008;39:1205–12.
- Saver JL, Jahan R, Levy EI, *et al*. Solitaire flow restoration device versus the Merci Retriever in patients with acute ischaemic stroke (SWIFT): a randomised, parallel-group, non-inferiority trial. *Lancet* 2012;380:1241–9.
- Nogueira RG, Lutsep HL, Gupta R, *et al*. Trevo versus Merci retrievers for thrombectomy revascularisation of large vessel occlusions in acute ischaemic stroke (TREVO 2): a randomised trial. *Lancet* 2012;380:1231–40.
- Zaidat OO, Castonguay AC, Gupta R, *et al*. North American Solitaire Stent Retriever Acute Stroke registry: post-marketing revascularization and clinical outcome results. *J Neurointerv Surg* 2014;6:584–8.
- No authors listed]. Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *N Engl J Med* 1995;333:1581–7.
- Rha JH, Saver JL. The impact of recanalization on ischemic stroke outcome: a meta-analysis. *Stroke* 2007;38:967–73.
- Wechsler LR, Roberts R, Furlan AJ, *et al*. Factors influencing outcome and treatment effect in PROACT II. *Stroke* 2003;34:1224–9.
- Singer OC, Haring HP, Trenkler J, *et al*. Age dependency of successful recanalization in anterior circulation stroke: the ENDOSTROKE study. *Cerebrovasc Dis* 2013;36:437–45.
- Ribo M, Flores A, Mansilla E, *et al*. Age-adjusted infarct volume threshold for good outcome after endovascular treatment. *J Neurointerv Surg* 2014;6:418–22.
- Yoo AJ, Chaudhry ZA, Nogueira RG, *et al*. Infarct volume is a pivotal biomarker after intra-arterial stroke therapy. *Stroke* 2012;43:1323–30.
- Saposnik G, Cote R, Phillips S, *et al*. Stroke outcome in those over 80: A multicenter cohort study across Canada. *Stroke* 2008;39:2310–17.
- Kaplan RC, Tirschwell DL, Longstreth WT Jr, *et al*. Vascular events, mortality, and preventive therapy following ischemic stroke in the elderly. *Neurology* 2005;65:835–42.
- Hussein HM, Georgiadis AL, Vazquez G, *et al*. Occurrence and predictors of futile recanalization following endovascular treatment among patients with acute ischemic stroke: a multicenter study. *AJNR Am J Neuroradiol* 2010;31:454–8.
- Sarraj A, Albright K, Barreto AD, *et al*. Optimizing prediction scores for poor outcome after intra-arterial therapy in anterior circulation acute ischemic stroke. *Stroke* 2013;44:3324–30.
- Flint AC, Xiang B, Gupta R, *et al*. THRIVE score predicts outcomes with a third-generation endovascular stroke treatment device in the TREVO-2 trial. *Stroke* 2013;44:3370–5.
- Ciccone A, Valvassori L, Nichelatti M, *et al*. Endovascular treatment for acute ischemic stroke. *New Engl J Med* 2013;368:904–13.
- Broderick JP, Palesch YY, Demchuk AM, *et al*. Endovascular therapy after intravenous t-PA versus t-PA alone for stroke. *New Engl J Med* 2013;368:893–903.
- Nogueira RG, Yoo AJ, Buonanno FS, *et al*. Endovascular approaches to acute stroke, Part 2: A comprehensive review of studies and trials. *AJNR Am J Neuroradiol* 2009;30:859–75.
- Kwak HS, Hwang SB, Jin GY, *et al*. Predictors of functional outcome after emergency carotid artery stenting and intra-arterial thrombolysis for treatment of acute stroke associated with obstruction of the proximal internal carotid artery and tandem downstream occlusion. *AJNR Am J Neuroradiol* 2013;34:841–6.
- Wolpert SM, Bruckmann H, Greenlee R, *et al*. Neuroradiologic evaluation of patients with acute stroke treated with recombinant tissue plasminogen activator. The rt-PA Acute Stroke Study Group. *AJNR Am J Neuroradiol* 1993;14:3–13.
- Linforte I, Linas RH, Selim M, *et al*. Clinical and vascular outcome in internal carotid artery versus middle cerebral artery occlusions after intravenous tissue plasminogen activator. *Stroke* 2002;33:2066–71.

- 25 Zaidat OO, Suarez JL, Santillan C, *et al.* Response to intra-arterial and combined intravenous and intra-arterial thrombolytic therapy in patients with distal internal carotid artery occlusion. *Stroke* 2002;33:1821–6.
- 26 Saver JL. Time is brain—quantified. *Stroke* 2006;37:263.
- 27 Wardlaw JM, Murray V, Berge E, *et al.* Recombinant tissue plasminogen activator for acute ischaemic stroke: an updated systematic review and meta-analysis. *Lancet* 2012;379:2364–72.
- 28 Saver JL, Fonarow GC, Smith EE, *et al.* Time to treatment with intravenous tissue plasminogen activator and outcome from acute ischemic stroke. *JAMA* 2013;309:2480–8.
- 29 Pereira VM, Gralla J, Davalos A, *et al.* Prospective, multicenter, single-arm study of mechanical thrombectomy using Solitaire flow restoration in acute ischemic stroke. *Stroke* 2013;44:2802–7.
- 30 Shi ZS, Liebeskind DS, Xiang B, *et al.* Predictors of functional dependence despite successful revascularization in large-vessel occlusion strokes. *Stroke* 2014;45:1977–84.
- 31 Pexman JH, Barber PA, Hill MD, *et al.* Use of the Alberta Stroke Program Early CT Score (ASPECTS) for assessing CT scans in patients with acute stroke. *AJNR Am J Neuroradiol* 2001;22:1534–42.

