When performing diagnostic angiography to evaluate the cervical carotid bifurcation, the catheter is placed inside the CCA to image both the ICA and ECA. In the vast majority of cases when only the ICA or ECA is visualized, the catheter has been unintentionally advanced distal to the bifurcation. This is easily remedied by withdrawing the catheter into the CCA. In the case of an absent CCA one will continue to image only the ICA or ECA (depending on which origin is catheterized) until the catheter is withdrawn into the IA. Failure to recognize these vessels originating from the IA may lead to the incorrect conclusion that one is occluded.

CONCLUSION

Congenital absence of the CCA is a rare finding that should be considered when a cervical angiogram shows only one carotid artery branch. This finding can alter the flow dynamics of the intracranial vasculature and potentially result in aneurysm formation.

Competing interests None.

Patient consent Obtained.

Ethics approval This study was conducted with the approval of the Cedars-Sinai.

Provenance and peer review Not commissioned; not externally peer reviewed.

REFERENCES

1. **Boyd JD.** Absence of the right common carotid artery. *J Anat* 1934;**68**:551–7.
2. **Woodruff WW III, Strunsky VP, Brown NJ.** Separate origins of the left internal and external carotid arteries directly from the aortic arch: duplex sonographic findings. *J Ultrasound Med* 1995;**14**:867–9.
3. **Monaco EA 3rd, Jankowitz BT, Tyler-Kabara EC, et al.** Incidental discovery of an absent right common carotid artery demonstrated by digital subtraction angiography and magnetic resonance angiography. *Klin Neuroradiol* 2009;**19**:227–9.
4. **Cerese A, Rubegni E, Tassi R, et al.** Absence of the right common carotid artery. *Surg Radiol Anat* 2009;**31**:815–17.
5. **Maybody M, Uszynski M, Morton E, et al.** Absence of the common carotid artery: a rare vascular anomaly. *AJNR Am J Neuroradiol* 2003;**24**:711–13.
6. **Kunishio K, Yamamoto Y, Sunami N, et al.** Agenesis of the left internal carotid artery, common carotid artery, and main trunk of the external carotid artery associated with multiple cerebral aneurysms. *Surg Neurol* 1987;**27**:177–81.
7. **Chen L, Liu JM, Zhou D.** Congenital absence of the right common carotid artery, internal carotid artery and external carotid artery associated with anterior communicating artery aneurysm: a rare case. *Neurol Sci* 2008;**29**:485–7.
8. **Cherici C.** Vincenzo Malacarne (1744-1816): a researcher in neurophysiology between anatomophysiology and electrical physiology of the human brain. *C R Biol* 2006;**329**:319–29.
9. **Malacarne V.** Osservazioni sopra alcune arterie del corpo umano nello stato preternaturale. *Osservatore Chir (Torino)* 1784;**2**:119.
10. **Kosinski J.** Absence of the right common carotid, division of the brachiocephalic trunk into the subclavian, external carotid and internal carotid. *Gaz Med Paris* 1868;**23**:65.
11. **Lie TA.** *Congenital anomalies of the carotid arteries.* Amsterdam: Excerpta Medica Foundation, 1968.
12. **Roberts LK, Gerald B.** Absence of both common carotid arteries. *AJR Am J Roentgenol* 1978;**130**:981–2.
13. **Rossitti S, Raininko R.** Absence of the common carotid artery in a patient with a persistent trigeminal artery variant. *Clin Radiol* 2001;**56**:79–81.
14. **Andrews ET, Howard JM.** Congenital anomalies of the carotid artery: a case report. *J Am Geriatr Soc* 1963;**11**:642–6.
15. **Halstuk KS, Littooy FN, Baker WH.** Absent common carotid artery associated with amaurosis fugax: a case report. *Surgery* 1985;**97**:502–6.
16. **Jerius JT, Stevens SL, Freeman MB, et al.** Vertebrobasilar syndrome associated with subclavian origin of the right internal carotid artery. *J Vasc Surg* 1995;**21**:855–61.
17. **Moncada R, Shannon M, Miller R, et al.** The cervical aortic arch. *Am J Roentgenol Radium Ther Nucl Med* 1975;**125**:591–601.
18. **Sadler TW.** *Langman's medical embryology.* 8th edn. Baltimore: Lippincott Williams & Wilkins, 2000.
19. **Pereira LP, Nepomuceno LA, Coimbra PP, et al.** Persistent trigeminal artery: angio-tomography and angio-magnetic resonance finding. *Arq Neuropsiquiatr* 2009;**67**:882–5.
20. **Eluvathingal Muttikkal TJ, Varghese SP, Chavan VN.** Persistent trigeminal artery and associated vascular variations. *Australas Radiol* 2007;**51** Spec No.:B31–3.
21. **Waaiajer A, van Leeuwen MS, van der Worp HB, et al.** Anatomic variations in the circle of Willis in patients with symptomatic carotid artery stenosis assessed with multidetector row CT angiography. *Cerebrovasc Dis* 2007;**23**:267–74.
22. **Chung TS, Lee YJ, Kang WS, et al.** Clinical and experimental investigation of pseudoaneurysm in the anterior communicating artery area on 3-dimensional time-of-flight cerebral magnetic resonance angiography. *J Comput Assist Tomogr* 2004;**28**:414–21.

Correction

The article “W S Lesley, R Rangaswamy. Catheter exchange technique for the Wingspan stent: technical note” was inadvertently published twice with different DOIs and should not be considered retracted. It was also published as part of the SNIS meeting abstracts in the July 2009 issue.

The correct citation is:

Lesley WS, Rangaswamy R. Catheter exchange technique for the Wingspan stent: technical note. *J NeuroIntervent Surg* 2009;**1**:168–70; doi:10.1136/jnis.2009.001867.

The journal apologises for the inadvertent duplicate publication and confirms that this article has not been retracted.

J NeuroIntervent Surg 2010;**2**:210. doi:10.1136/jnis.2010.307corr2