performing MCI. Further prospective studies are warranted to evaluate if the MCI technique affects reperfusion and functional outcomes in endovascular patients.

Disclosures M. Nouri: None. J. Lee: None. J. Mocco: None. P. Singh: 6; C; Penumbra.

E-138 ENDOVASCULAR MANAGEMENT OF FREE FLOATING COMMON CAROTID THROMBI

¹K Carr*, ²Z Thwing, ²R Shivashakar, ²Y Serulle. ¹Aventura Hospital and Medical Center, Coral Springs, FL; ²Aventura Hospital and Medical Center, Aventura, FL

10.1136/neurintsurg-2019-SNIS.213

Background Free floating thrombi of the common carotid artery are rare causes of acute stroke syndromes. In previously described cases, authors have either advocated surgical management or conservative anticoagulation therapy. We described two cases in which endovascular management provided definitive and longstanding treatment.

Case 1 A 51-year-old Hispanic woman with a history of insulin dependent diabetes mellitus (IDDM), hypertension and hyperlipidemia presented with 2–3 days of waxing and waning right sided hemiparesis, aphasia and paresthesia. Initial non-contrast-enhanced head computed tomography (NECT) examination demonstrated no evidence of acute intracranial pathology. Doppler sonogram and cervical angiogram confirmed the presence of the thrombus with 70–80% stenosis. CT angiogram of the brain vasculature demonstrated no evidence of intracranial large vessel occlusion. Brain Magnetic Resonance Imaging (MRI) demonstrated acute watershed infarcts in the left cerebral hemisphere as well as a few punctate embolic infarcts. Serologic workup was significant only for hyperglycemia. After a brief course of heparinization, successful mechanical revascularization was accomplished using direct aspiration of the thrombus.

Case 2 A 51-year-old African American male with a history of insulin dependent diabetes mellitus and hyperlipidemia presented emergently after 6 hours of acute aphasia and right sided hemiparesis. Initial NECT demonstrated left middle cerebral artery (MCA) territory infarcts involving the frontal and temporal lobes. Cervical CT Angiogram demonstrated a long segment free floating thrombus of the distal left common carotid artery (CCA), which was confirmed on subsequent cervical angiogram. In this case, mechanical revascularization was accomplished with aspiration and stent retriever, followed by balloon angioplasty.

Conclusion Endovascular mechanical revascularization is an efficient and definitive treatment in patients with free floating carotid thrombi.

Disclosures K. Carr: None. Z. Thwing: None. R. Shivashakar: None. Y. Serulle: None.

E-139 THE CHARACTERISTICS OF ELDERLY PATIENTS OVER 80 YEARS OF AGE WHO UNDERWENT MECHANICAL THROMBECTOMY USING STENT-RETRIEVERS DUE TO ACUTE CEREBRAL INFARCTION

K Jang*, B Cho, B Moon, K Kim, D Jang. *Neurosurgery, Incheon St. Mary's hospital, The Catholic University of Korea, Bupyeong-gu, Incheon, Korea, republic of*

10.1136/neurintsurg-2019-SNIS.214

Introduction Recently, the average life expectancy of people has increased, and the number of cerebral infarctions in elderly patients over 80 years of age is also increasing. There are increasing cases of mechanical thrombectomy using stent-retriever in elderly patient, but the characteristics and differences of the patient population are not well established yet. The purpose of this study is to investigate the characteristics and prognostic factors of elderly patients over 80 years of age undergoing stent-retriever thrombectomy.

Methods From 2013 to 2017, we analyzed the prospective gathered data of 190 consecutive patients treated with mechanical thrombectomy using stent-retrievers for acute ischemic stroke in a single center. Demographic, clinical, laboratory, and radiologic features of each patient were investigated. We compared the characteristics of elderly patients over 80 year-old with those of younger patients and evaluated the prognostic factors associated with good outcomes in elderly patients.

Results Of the total 133 patients, 34 patients (25.6%) were over 80 years old. Initial ASPECT score (9.41 vs 8.78, $p=0.016$), 30 days mRS (4.21 vs 3.18, $p=0.003$) and 90 days mRS (4.15 vs 2.95, $p=0.001$) were significantly higher at elderly patients. Hypertension history ($p<0.001$), underlying atrial fibrillation ($p=0.007$) were significantly more common in elderly patients. The favorable outcomes ($mRS\leq 2$, $p=0.035$) were significantly different between the two groups. Because the 30 days and 90 days mRS were higher in the elderly patients after mechanical thrombectomy, prognosis may be worse and other complications should be noted.

Conclusions Elderly patients over 80 year-old had a few more underlying diseases, and the functional outcome was slightly worse after mechanical thrombectomy using stent retriever in acute cerebral infarction. However, the prognostic factors such as mortality and favorable outcome did not differ significantly between the two groups. Elderly patients should also consider aggressive treatment while appropriately controlling underlying diseases.

Disclosures K. Jang: None. B. Cho: None. B. Moon: None. K. Kim: None. D. Jang: None.

E-140 STRATEGIZING TANDEM OCCLUSION ACUTE STROKE MANAGEMENT BASED ON CTA

¹S Paramasivam*, ²S Kumar. ¹Neurosurgery, Apollo Hospitals, Chennai, India; ²Neurosurgery, Apollo Hospitals, Bangalore, India

10.1136/neurintsurg-2019-SNIS.215

Objectives To strategize the management of major stroke secondary to tandem occlusions of extracranial ICA and Intracranial ICA/MCA based on the collateral assessment in the pre-procedure CTA.

Methods In a retrospective analysis of our Stroke Thrombectomy database ($n = 85$), treated between August 2016 to December 2018 we identified 15 patients who presented with tandem occlusion. 13 were atherosclerotic disease and 2 were carotid dissection along with middle cerebral artery or distal ICA occlusion. All patients had pre-procedure CT angiogram, along with the primary pathology, collaterals comprising of Acom and Pcom arteries and their caliber was analyzed. All except 3 patients were treated with balloon