

SUPPLEMENTARY MATERIAL

Supplementary table 1: Demographic baseline characteristics of the ESCAPE ALICE participants (N = 248).

Gender – n (%)	37 (14.9) females/ 211 (85.1) males
Age – n (%)	
21 – 30	6 (2.4)
31 – 40	75 (30.2)
41 – 50	111 (44.8)
51 – 60	43 (17.3)
61 – 70	13 (5.2)
Specialty – n (%)	
Neurointerventional Radiology	193 (77.8)
Neurosurgery*	36 (14.5)
Neurology*	12 (4.8)
Other	7 (2.8)
Career stage – n (%)	
Resident	6 (2.4)
Fellow	3 (1.2)
Junior Staff†	37 (14.9)
Senior Staff‡	121 (48.8)
Division/Department Head	81 (32.7)
Neurovascular experience in years – n (%)	
Less than 5	50 (20.2)
5 – 10	61 (24.6)
10 – 15	68 (27.4)
15 – 20	32 (12.9)
More than 20	37 (14.9)
Personal annual thrombectomy volume – n (%)	
Less than 10	31 (12.6)
10 - 50	133 (53.9)
50 - 100	69 (27.9)
100 - 150	10 (4.1)
150 - 200	2 (0.8)
More than 200	2 (0.8)
Institutional annual thrombectomy volume – n (%)	
Less than 10	14 (5.7)
10 - 50	52 (21.1)
50 - 100	45 (18.2)
100 - 150	47 (19.0)
150 - 200	31 (12.6)
200 - 250	25 (10.1)
250 - 300	12 (4.9)
More than 300	21 (8.5)
Hospital catchment population – n (%)	
Less than 1 million	75 (30.2)
1 – 2 million	107 (43.1)
2 – 3 million	29 (11.7)
3 – 4 million	13 (5.2)
4 – 5 million	8 (3.2)
More than 5 million	18 (5.2)
Not sure	3 (1.2)
Number of neurointerventionalists taking calls – n (%)	
1 - 3	121 (48.8)
4 - 6	114 (46.0)
7 - 9	10 (4.0)
10 or more	3 (1.2)

* All neurologists/neurosurgeons were endovascularly trained

† Less than 5 years from board certification

‡ More than 5 years from board certification

Supplementary figure 1: Location of the Emergency Department in relation to the CT scanner (left) and of the CT scanner in relation to the angiography suite (right) in survey participants' institutions. Dark blue color indicates location on the same floor nearby, orange color indicates location on the same floor but remote, and light blue indicates location on a different floor. Red indicates a hybrid setup/setup in which the CT scanner is bypassed by default.