Conclusion According to our results, CA in non-complicated cases can be safely performed outside of hospital setting by board-certified interventional neuroradiologists with high success rates and minimal complications that are comparable to MRA and CTA. Our result show that not only CA is safe in an office setting, but potentially safer than MRA and CTA when considering higher quality imaging, evaluation of collateral flow and diagnosis of many additional vascular diseases. Proper case selection for this setting plays an important role in achieving optimal results and minimizing complications.

Disclosures S. Razavi: None. E. Masangkay: None. N. Chelikam: None. U. Kelly-Tolley: None. L. Pierce: None. R. Malek: None. A. Padiar: None.

E-159 FEASIBILITY AND SAFETY OF TRANSRADIAL ACCESS FOR PEDIATRIC NEUROINTERVENTIONS

¹V Srinivasan*, C Hadley, 77030¹, ¹M Prabek, ¹M Lopresti, ²S Chen, ²E Peterson, ³A Sweid, ³P Jabbar, ⁴C Young, ⁴M Levitt, ⁵J Osbun, ¹J Burkhardt, ¹J Johnson, ¹P Kan. ¹Neurosurgery, Baylor College of Medicine, Houston, TX; ²Neurosurgery, University of Miami, Miami, FL; ³Neurosurgery, Thomas Jefferson University, Philadelphia, PA; ⁴Neurosurgery, University of Washington, Seattle, WA; ⁵Neurosurgery, Washington University School of Medicine, St. Louis, MO

10.1136/neurintsurg-2020-SNIS.191

Background Diagnostic cerebral angiograms are increasingly being performed by transradial access (TRA) in adults, following in line with data from the coronary literature supporting fewer access-site complications. Despite this ongoing trend on TRA in neuroangiography, there has been no discussion of its use in the pediatric population. In fact, pediatric TRA for other endovascular intervention has scarcely been described even for coronary or other applications. This is the first dedicated study of transradial access for neuroangiography in pediatric patients.

Methods A multi-institutional series of consecutively performed pediatric transradial angiograms and interventions was collected. This included demographic, procedural, outcomes, and safety data. Data was prospectively recorded and retrospectively analyzed.

Results A total of 37 diagnostic angiograms and 24 interventions were performed in 47 pediatric patients, by 5 neurointerventionalists. Proximal and distal angiography were performed successfully for both diagnostic and interventional application (19 distal angiograms, 2 distal interventions). Clinically significant vasospasm occurred in 7 patients (11.5%). Re-access was successfully performed in 7 patients a total of 11 times. Conversion to femoral access occurred in 6 cases (9.8%). There were no access-related complications otherwise.

Conclusions Transradial access in pediatric patients is safe and feasible. It can be performed successfully in many cases but carries some unique challenges compared to the adult population. Despite the challenge of higher rates of vasospasm and conversion to transfemoral access, it is worth exploring further, given the potential benefits.

Disclosures V. Srinivasan: None. C. Hadley: None. M. Prabek: None. M. Lopresti: None. S. Chen: None. E. Peterson: None. A. Sweid: None. P. Jabbar: None. C. Young: None. M. Levitt: None. J. Osbun: None. J. Burkhardt: None. J. Johnson: None. P. Kan: None.

E-160 BLOOD CLOT IDENTIFICATION AND COMPOSITION ASSESSMENT BY FAST SPIN-ECHO (FSE) T2WI AND T2* MAPPING

Y Ding*, M Abbassi, J Felmlee, D Dai, R Kadirvel, D Kallmes, W Brinjikji. *Radiology, Mayo Clinic, Rochester, MN*

10.1136/neurintsurg-2020-SNIS.192

Purpose There are growing data to suggest that clot composition can impact revascularization outcomes and strategies in large vessel occlusion patients. In many centers, MRI is the primary modality of acute stroke imaging so identifying clot composition on MRI imaging may be important. We performed an in vitro study to determine the MR signaling characteristics of stroke clots of various compositions.

Methods Fifteen thrombus analogs of five compositions (n=3 for each composition) [Group A, fibrin-rich (95% plasma:5% RBCs); Group B, fibrin-rich (75% plasma:25% RBCs); Group C, intermediate (50% plasma:50% RBCs); Group D, RBC-rich (25% plasma:75% RBCs,) and Group E, RBC-rich (5% plasma:95% RBCs)] were scanned with fast spin-echo (FSE) T2WI (TR/TE 2500/101 milliseconds (ms)) and quantitative T2* mapping sequence. Signals from FSE T2WI were collected, and thrombus T2* relaxation time (TT2*RT) was measured in all the groups. Correlation between the thrombus-T2* relaxation time and red blood cell content was analyzed.

Results Signal intensity changed gradually from high (bright) to low (dark) from Group A to E gradually from all the 3 clots in each group, which indicated the signal intensity was decreased as the composition of RBC increased. The average TT2*RT decreased from 60 ms (green) to 25 ms (red) from fibrin-rich clot to RBC-rich clot (Group A to E), which

Abstract E-160 Figure 1