(OR=0.544, [95% CI 0.4- 0.7]; $p=0.032$), which indicates that patients with robust VMP had better neurological outcomes 90 days after discharge.

Conclusion A robust cerebral venous microperfusion profile reflects greater tissue microperfusion, good arterial collateralization status and is associated with improved clinical outcome in patients with AIS.

Disclosures T. Faizy: None. R. Kabiri: None. M. Leipzig: None. S. Christensen: None. G. Broocks: None. F. Flottmann: None. M. Lansberg: None. G. Albers: None. J. Fiehler: None. M. Wintermark: None. J. Heit: None.

O-003

RESULTS OF THE STROKE THROMBOEMBOLISM REGISTRY OF IMAGING AND PATHOLOGY: A MULTICENTER INTERNATIONAL STUDY

¹W Brinjikji*, ²R Nogueira, ³R Hanel, ⁴K Layton, ⁵J Delgado, ⁶M Gounis, ⁷A Yoo, ⁸V Pereira, ⁹M Almekhlafi, ¹⁰K Doyle, ¹¹B Patel, ¹²B Jahromi, ¹³P Kvamme, ¹R Kadirvel, ¹D Dai, ¹D Kallmes, ¹S Fitzgerald. ¹Radiology, Mayo Clinic, Rochester, MN; ²Neurology, Grady Memorial Hospital, Atlanta, GA; ³Neurosurgery, Baptist, Jacksonville, FL; ⁴Radiology, Baylor Scott White Hospital, Dallas, TX; ⁵Radiology, Abbott Northwestern Hospital, Minneapolis, MN; ⁶Radiology, University of Massachusetts, Worcester, MA; ⁷Radiology, UT Houston, Houston, TX; ⁸Neurosurgery, University of Toronto, Toronto, ON, CANADA; ⁹Neurology, University of Calgary, Calgary, AB, CANADA; ¹⁰Physiology, University of Galway, Galway, IRELAND; ¹¹Radiology, Carilion Clinic, Carilion, WV; ¹²Neurosurgery, Northwestern Memorial Hospital, Chicago, IL; ¹³Radiology, UT Knoxville, Knoxville, TN

10.1136/neurintsurg-2020-SNIS.3

Background We performed a multicenter prospective clinical registry across 12 centers to study the association between histopathological characteristics of retrieved clots and imaging, stroke etiology and clinical outcomes.

Materials and Methods Following IRB approval at the 12 centers, patients were enrolled in the STRIP registry. All retrieved emboli were sent for histopathological analysis with H&E and MSB staining. Demographic variables, comorbidities, stroke etiology, imaging findings and procedural details were collected for each case. We studied the association between clot histopathology and imaging findings, stroke etiology and and revascularization outcomes. Student's t-test was used for continuous variables and chi-squared testing for categorical variables.

Results To date, 1457 patients have been included. Platelet rich clots were associated with a significantly lower rate of first pass TICI 2c/3 revascularization than platelet poor clots (OR=0.68, 95%CI=0.48–0.88, $P<0.0001$). The rate of first pass TICI 2c/3 recanalization was 41% with aspiration compared to 26% for stent-trieter and 15% for combined

techniques with platelet rich clots ($P<0.0001$). There was a significant correlation between platelet rich clots and the absence of hyperdensity on non-contrast CT ($p=0.003$) and a significant inverse correlation between the percentage of platelets and mean HU on NCCT ($r=-0.243$, $p=0.005$). We found that a Hounsfield unit cutoff of 55 or less had a sensitivity and specificity of 70% for a platelet rich clot. We did not notice any interaction between clot composition and stroke etiology.

Conclusions Interim analysis of the STRIP registry suggests that the platelet content of a clot may be the most revealing factor in determining a clot's imaging features and revascularization outcome. Platelet rich clots are less dense on NCCT and are associated with poorer first pass TICI 2c/3. Aspiration therapy was associated with higher rates of first pass recanalization than stent-retriever thrombectomy or combined techniques.

Disclosures W. Brinjikji: None. R. Nogueira: None. R. Hanel: None. K. Layton: None. J. Delgado: None. M. Gounis: None. A. Yoo: None. V. Pereira: None. M. Almekhlafi: None. K. Doyle: None. B. Patel: None. B. Jahromi: None. P. Kvamme: None. R. Kadirvel: None. D. Dai: None. D. Kallmes: None. S. Fitzgerald: None.

O-004

PROSPECTIVE, MULTI-CENTERED, EMS-ADMINISTERED, PRE-HOSPITAL VALIDATION STUDY OF THE RAPID ARTERIAL OCCLUSION EVALUATION (RACE) SCALE FOR DETECTING LARGE VESSEL OCCLUSION STROKE IN THE UNITED STATES COMPARED TO THE ORIGINAL RACE VALIDATION STUDY FROM SPAIN: A SUBANALYSIS OF THE PREDICT STUDY

¹A Cruz*, ¹E Fortuny, ¹B Ugiliweneza, ¹D Wang, ²A White, ¹N Khattar, ¹S Adams, ¹B Gallinore, ¹D Ding, ³S Wolfe, ⁴D Heck, ¹R James. ¹Neurosurgery, University of Louisville, Louisville, KY; ²Radiology, University of Louisville, Louisville, KY; ³Neurosurgery, Wake Forest Baptist Health, Lexington, NC; ⁴Radiology, Forsythe Medical Center, Winston-Salem, NC

10.1136/neurintsurg-2020-SNIS.4

Introduction Pre-hospital identification of patients with large vessel occlusion (LVO) is critical in the timely triage of thrombectomy-eligible LVO patients to comprehensive stroke centers. The Rapid Arterial occlusion Evaluation (RACE) scale was prospectively validated to identify LVO patients by EMS in Spain and has been widely adopted in the United States, though these healthcare infrastructure and EMS systems may differ significantly and these results may not be completely

Abstract O-004 Table 1 Sensitivity, Specificity, Accuracy, PPV and NPV values for the RACE scale across various cut-off values for PREDICT (U.S based study) and the Original RACE study (Spain Based) PPV=Positive Predictive Value, Negative Predictive Value

RACE Score	RACE Scale (U.S Based study)					RACE scale (Original Spanish Based Study)				
	Sensitivity	Specificity	PPV	NPV	Accuracy	Sensitivity	Specificity	PPV	NPV	Accuracy
≥1	1.00	0.149	0.119	1.00	0.237	1.00	0.13	0.24	1.00	0.31
≥2	0.958	0.303	0.137	0.984	0.371	0.97	0.27	0.27	0.97	0.42
≥3	0.875	0.519	0.174	0.973	0.556	0.93	0.40	0.30	0.96	0.51
≥4	0.833	0.625	0.204	0.970	0.647	0.89	0.55	0.35	0.95	0.62
≥5	0.792	0.716	0.244	0.968	0.724	0.85	0.68	0.42	0.94	0.72
≥6	0.750	0.784	0.286	0.964	0.780	0.72	0.77	0.46	0.91	0.76
≥7	0.500	0.865	0.300	0.938	0.828	0.53	0.89	0.56	0.87	0.81
≥8	0.250	0.947	0.353	0.916	0.875	0.32	0.95	0.65	0.84	0.82
9	0.167	0.986	0.571	0.911	0.901	0.07	0.99	0.56	0.79	0.79

applicable in the US. This study prospectively validates the RACE scale as administered by US-based EMS personnel in the pre-hospital setting and we compare our results to the originally published results.

Methods 232 adult patients suspected of having a stroke by EMS and transported to a participating comprehensive stroke center had the RACE scale administered prospectively and recorded in a secure web-based database. Admission NIHSS score and final diagnosis were recorded. Cerebrovascular imaging studies (CTA, MRA or DSA) were reviewed by a blinded, independent Neuroradiologist to determine LVO diagnosis. We used SAS and c-statistics to create receiver operating characteristic (ROC) curves to determine the area under the curve (AUC) and optimal cut point (CP) scores for the RACE scale. We also calculated the sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), overall accuracy for the optimal CP score.

Results Our analysis of the predictive capability of the RACE scale showed similar predictive capacities for radiographically-confirmed LVO in patients prospectively tested in the US as compared to the original Spanish RACE scale population. The best CP score value predictive of LVO in our US-RACE study was determined to be ≥ 6 , compared to the original Spanish study which was ≥ 5 (table 1). The overall prevalence of LVO as defined in our study was 13.4%, compared to a prevalence of 21.3% for the original paper's definition of LVO.

Conclusion This is the first prospective validation of the RACE scale performed in the US. These results demonstrate that the RACE scale retains the previously published predictive value in both the US and Spain in accurately identifying LVO stroke in a prehospital setting by EMS.

Disclosures A. Cruz: None. E. Fortuny: None. B. Ugiliweneza: None. D. Wang: None. A. White: None. N. Khattar: None. S. Adams: None. B. Gallimore: None. D. Ding: None. S. Wolfe: None. D. Heck: None. R. James: None.

O-005 VASCULAR COLLATERALIZATION MAY NOT AFFECT BLOOD GAS CHANGES IN PERI-INFARCT VASCULATURE IN HUMAN ISCHEMIC STROKE

¹R Spears*, ²C McLouth, ³K Pennypacker, ¹J Frank, ⁴B Maglinger, ⁴A Trout, ⁴J Roberts, ⁵A Stowe, ¹S Grupke, ¹J Fraser. ¹Neurosurgery, University of Kentucky, Lexington, KY; ²Behavioral Science, University of Kentucky, Lexington, KY; ³Neurology and Anatomy, University of Kentucky, Lexington, KY; ⁴Neuroscience, University of Kentucky, Lexington, KY; ⁵Neurology, University of Kentucky, Lexington, KY

10.1136/neurintsurg-2020-SNIS.5

Introduction Ischemic stroke is a prevalent, devastating disease with high morbidity and mortality. Despite extensive research using animal models, there remains significant gaps in understanding processes of stroke in human patients. To address this, we developed a protocol to obtain and to analyze blood immediately proximal in systemic circulation and distal to a thrombus in patients undergoing mechanical thrombectomy (www.clinicaltrials.gov NCT03153683). Our goal for this project was to evaluate blood gas changes and acid/base balance during stroke and how these changes are affected by patient factors.

Methods We analyzed blood samples from the first 62 patients in the BACTRAC registry. Bicarbonate, pO₂, and pCO₂ values of intracranial (distal) and systemic (proximal) arterial blood relative to the occlusive thrombus were analyzed. Changes

Abstract O-005 Table 1 Comparison by CTA score of systemic and intracranial blood gasses

	CTA 0 (n=17)	CTA 1+ (n=41)	p-value
Systemic			
pO ₂	211.35 (64.08)	255.22 (93.98)	0.084
pCO ₂	39.43 (5.19)	37.66 (4.48)	0.196
Bicarbonate	22.38 (2.62)	22.12 (3.47)	0.787
pH	7.36 (0.04)	7.38 (0.06)	0.407
Intracranial			
pO ₂	182.25 (48.83)	215.69 (66.44)	0.077
pCO ₂	33.20 (9.83)	31.29 (8.87)	0.492
Bicarbonate	18.91 (5.97)	18.39 (5.61)	0.766
pH	7.36 (0.04)	7.37 (0.07)	0.426

were compared in patients according to vascular collateralization as measured by CTA collateral scores.

Results Mean age was 68.9 years (25 – 95 years). 29 were male, 33 were female. 15 were current smokers (24%), and 47 were non-smokers (no smoking within the last 6 months; 76%). Overall, intracranial gas values differed significantly from systemic. Compared to systemic, mean intracranial pO₂ was decreased (211.39 vs. 246.91, $p < 0.001$), pCO₂ was decreased (32.19 vs. 38.12, $p < 0.001$, and bicarbonate was decreased (18.90 vs. 22.20, $p < 0.001$). Collateralization did not significantly affect distal blood gas values.

Discussion and Conclusion A compensated metabolic acidosis is present in arterial blood gas samples immediately proximal and distal to thrombi in large vessel occlusive stroke. Vascular collateralization may not significantly affect the acid-base environment immediately distal to a large vessel occlusion.

Disclosures R. Spears: None. C. McLouth: None. K. Pennypacker: None. J. Frank: None. B. Maglinger: None. A. Trout: None. J. Roberts: None. A. Stowe: None. S. Grupke: None. J. Fraser: None.

O-006 HISTOTRIPSY FOR INTRACEREBRAL HEMORRHAGE IN A PORCINE MODEL

T Gerhardson, J Sukovich, N Chaudhary, T Chenevert, K Ives, T Hall, S Camelo-Piragua, B Daou*, Z Xu, A Pandey. Neurosurgery, University of Michigan, Ann Arbor, MI

10.1136/neurintsurg-2020-SNIS.6

Introduction Histotripsy is a noninvasive, focused ultrasound technique that generates cavitation to mechanically fractionate tissue. Intracerebral hemorrhage (ICH) is characterized by a 30-d mortality rate of 40% and significant disability for those who survive. Histotripsy has the potential to liquefy clot in the brain and facilitate minimally invasive aspiration. We aim to investigate the initial safety concerns of histotripsy mediated clot liquefaction and aspiration in a porcine ICH model.

Methods About 1.75-mL clots were formed in the frontal lobe of the brain (n=18; n=6/group). The centers of the clots were liquefied with histotripsy 48 h after formation, and the content was either evacuated or left within the brain. A control group was left untreated. Pigs underwent magnetic resonance imaging (MRI) 7 to 8 d after clot formation and were subsequently euthanized. Neurological behavior was assessed throughout. Histological analysis was performed on harvested brains. A subset of pigs underwent acute analysis (≤ 6 h).