Grande: None. **R. Crosa:** None. **S. Webb:** None. **M. Psychios:** None. **E. Peterson:** None. **D. Yavagal:** None. **A. Spiotta:** 1; C; Penumbra, Medtronic, Stryker. 2; C; Cerenovus, Terumo, Siemens, Penumbra, Stryker. **R. Starke:** 1; C; NREF, Joe Niekro Foundation, Brain Aneurysm Foundation, Bee Foundation, NIH, Miami Clinical and Translational Science Institute, National Center for Advancing Translational Sciences, National Institute on Minority Health and Health Disparities. 2; C; Penumbra, Abbott, Medtronic, InNeuroCo, Cerenovus.

P-009

AN ANGLED TIP IS ASSOCIATED WITH IMPROVED TECHNICAL OUTCOMES WHEN USING 0.064–0.074 ASPIRATION CATHETERS: ANALYSIS FROM A MULTICENTER RETROSPECTIVE COHORT

¹J Vargas*, ²S Majidi, ³H Hawk, ⁴S Nimjee, ⁵A Zakeri, ⁶M Mokin, ⁷R Kellogg, ⁸G Cortez, ⁹A Aghaebrahim, ¹⁰E Sauvageau, ¹¹R Hanel, ¹²R De Leacy, ¹³A Siddiqui, ¹⁴A Turk, ¹⁵M Oselkin, ¹⁶E Marlin, ¹⁷R Turner, ¹⁸I Chaudry, ¹⁹J Milburn. ¹Neurosurgery, Southeastern Neurosurgery and Spine, Greenville, SC; ²Neurosurgery, Mount Sinai Hospital, New York, NY; ³Radiology, Erlanger Health System, Chattanooga, TN; ⁴Neurosurgery, The Ohio State University Medical Center, Columbus, OH; ⁵Neurosurgery, Tampa General Hospital, Tampa, FL; ⁶Neurosurgery, University of Virginia, Charlottesville, VA; ⁷Neurosurgery, Baptist MC, Jacksonville, FL; ⁸Neurosurgery, Baptist Health, Jacksonville, FL; ⁹Neurosurgery, Mount Sinai, New York, NY; ¹⁰Neurosurgery, University of Buffalo, Buffalo, NY; ¹¹Neurosurgery, MUSC, Greenville, SC; ¹²Radiology, St Luke's University, Bethlehem, PA; ¹³Neurosurgery, St Luke's University, Bethlehem, PA; ¹⁴Neurosurgery, Prisma Health, Greenville, SC; ¹⁵Radiology, Ochsner Health, New Orleans, LA

10.1136/neurintsurg-2021-SNIS.45

Background Recent introduction of large bore aspiration catheters have contributed to high reperfusion rates. The alteration of these catheters to have angled tips, which increase clot engagement area by approximately 15%, has been suggested to improve rates of complete clot ingestion, resulting in higher quality thrombectomies. We aimed to evaluate the recanalization efficacy of new generation angled tip aspiration catheters in comparison to commonly used straight tip aspiration catheters in real world data set.

Methods We performed a multicenter retrospective analysis of consecutive acute ischemic stroke patients with M1 occlusion treated within 24 hours from the time of last known well. Patients were divided into two cohorts: those in whom a 0.071 angled tip catheter was the initial device used to attempt reperfusion and those in whom a 0.064–0.074 inch flat tip (non-angled) catheter was the initial device used to attempt reperfusion. The primary outcome was the rate of excellent reperfusion (TICI 2C). Secondary outcomes included the rate of successful reperfusion (TICI2B), use of stent retrievers, and access to successful reperfusion time. All data was self-adjudicated. No outside funding was provided for this analysis.

Results Total of 650 patients with acute M1 occlusion who underwent thrombectomy were identified. Angled tip Zoom 71 catheter was used in 162 patients, while 488 patients were treated with flat tip aspiration catheters (ranging 0.064 to 0.074 inch in inner diameter). The baseline mRS score, admission NIHSS score, the rate of intravenous thrombolytic therapy, symptom onset to hospital arrival, and use of anesthesia were not different between the cohorts. The primary outcome was significantly higher in the angled tip cohort (67% vs 59%, $p=0.03$). There was no difference in the rate of TICI2B reperfusion (96% vs 94%, $p=0.71$). There was a lower rate of stent retriever use (25% vs 44%, $p<0.0001$) in the angled

tip cohort. Access time to successful reperfusion was significantly faster with angled tip catheters (21 vs 29 minutes, $p<0.0001$). Access time to TICI 2B (25 vs 31 minutes, $p=0.03$) and final recanalization (29 vs 35 minutes, $p=0.03$) remained significantly shorter in the angled tip cohort, after adjusting for age, thrombectomy technique, use of secondary aspiration catheter.

Conclusion This multicenter, consecutive real-world experience demonstrates that M1 thrombectomy with an angled tip aspiration catheter is associated with higher rates of TICI 2C or better reperfusion, equal rates of TICI 2B or better reperfusion, and faster time to successful reperfusion.

Disclosures J. Vargas: None. S. Majidi: None. H. Hawk: None. S. Nimjee: None. A. Zakeri: None. M. Mokin: None. R. Kellogg: None. G. Cortez: None. A. Aghaebrahim: None. E. Sauvageau: None. R. Hanel: None. R. De Leacy: None. A. Siddiqui: None. A. Turk: None. M. Oselkin: None. E. Marlin: None. R. Turner: None. I. Chaudry: None. J. Milburn: None.

P-010

FIRST LINE DIRECT CONTACT ASPIRATION VERSUS STENT-RETRIEVER BASED TECHNIQUES: REAL-WORLD COMPARATIVE THROMBECTOMY OUTCOMES FROM THE NVQI-QOD ACUTE ISCHEMIC STROKE REGISTRY

¹A Mahapatra*, ²Y Moazeni, ³T Patterson, ⁴R Abdalla, ⁵J Tsai, ⁶N Moore, ⁷M Bain, ⁸G Toth, ⁹S Ansari, ¹⁰M Hussain. ¹Cerebrovascular Center, Cleveland Clinic Foundation, Cleveland, OH; ²Radiology, Northwestern University, Chicago, IL; ³Neurology, Spectrum Health, Grand Rapids, MI

10.1136/neurintsurg-2021-SNIS.46

Introduction Mechanical thrombectomy is the standard of care treatment for large-vessel acute ischemic stroke and associated with significant improvement in long-term functional outcomes. Early and complete recanalization are paramount in achieving optimal patient outcomes. Though several revascularization techniques have been described, direct contact aspiration and clot removal via stent-retriever remain the foundation of endovascular stroke thrombectomy. Utilizing the NeuroVascular Quality Initiative - Quality Outcomes Database (NVQI-QOD) Acute Ischemic Stroke Registry, we present real-world data on first-line practice for treatment of emergent large vessel occlusions (ELVOs), comparing angiographic and clinical outcomes between direct contact aspiration and stent-retriever mechanical thrombectomy techniques.

Methods Retrospective analysis of the NVQI-QOD registry data was performed. We included patients with ELVOs that underwent mechanical thrombectomy with age > 18 years and presenting NIHSS ≥ 6 . We compared procedural times, recanalization efficacy, and clinical outcomes, including in-hospital mortality and discharge NIHSS.

Results We identified a total of 2381 patients that met inclusion criteria. 998 (41.9%) underwent treatment with direct contact aspiration alone and 1383 (58.1%) underwent treatment utilizing a stent-retriever (with or without local aspiration). There were no significant differences in baseline median NIHSS scores (16 vs. 17, $p = 0.25$) or median ASPECTS scores (9 vs. 9, $p = 0.7$). No significant differences were seen in time last known well to puncture times (282 min vs. 280 min, $p = 0.22$) or recanalization time (323 min vs. 322 min, $p = 0.39$), ED to puncture time (75 min vs. 71 min, $p = 0.25$) or recanalization time (158 min vs. 160 min, $p = 0.55$), or median procedure times between the two groups (23