negative linear correlation with the number of new or cumulative COVID-19 cases in Hungary.

Conclusions Collateral damages of the COVID-19 outbreak on the Hungarian IS care system were demonstrated. The disproportionately greater decline of IS admission numbers could partially be explained by the effect of health emergency operative measures. Further studies are needed to evaluate the causes of our observations.

REFERENCES

1. Bersano A, Kraemer M, Touzé E, Weber R, Alamowitch S, Sibon I, et al. Stroke care during the COVID-19 pandemic: experience from three large European countries. *Eur J Neurol* 2020;27(9):1794–800. <https://doi.org/10.1111/ene.14375>
2. Böjti PP, Stang R, Gunda B, Sipos I, Bereczki D. A járulékos egészségügyi veszteségek retrospektív, egycentrumos felmérése. [Effects of COVID-19 pandemic on acute ischemic stroke care. A single-centre retrospective analysis of medical collateral damage]. *Orv Hetil* 2020;161(34):1395–9. <https://doi.org/10.1556/650.2020.31936>. Hungarian.
3. Slowik A, Nowak R, Popiela T. Significant fall in stroke admissions in the Malopolska Voivodeship of Poland during the COVID-19 pandemic. *Neurol Neurochir Pol* 2020;54(5):471–2. <https://doi.org/10.5603/PJNNS.a2020.0056>

Disclosure Nothing to disclose

EP40

TELEPROCTORING IN NEUROINTERVENTIONISM

B Pabon, V Torres, M Patiño, N Serna, O Vargas, C Estrada, J Pelaez, J Mejia. *Angioteam, Medellín, Colombia*

10.1136/neurintsurg-2021-ESMINT.39

6 cases of neurovascular treatment with remote assistance are presented. various techniques (zoom, smart glasses and hospital internal system) 2 aneurysm treatments with flow diversion are presented. 1 case with web device. and 3 cases of stroke.

All cases occurred during the covid 19 pandemic.

Multiple tests were carried out during the simulation courses. Evaluating communication in real time. mastery of the images and mastery of the software.

REFERENCE

1. Bechstein M, et al. Interhospital teleproctoring of endovascular intracranial aneurysm treatment using a dedicated live-streaming technology: first experiences during the COVID-19 pandemic. *BMJ Case Rep* 2020. PMID: 33012707.

Disclosure Boris Pabon: proctorship con MEDTRONIC, Microvention Consultant MIVI

Imaging

EP41*

WHICH TRANSFER PATIENTS WITH STROKE DUE TO A LARGE-VESSEL OCCLUSION BENEFIT FROM CT SCAN REEVALUATION PRIOR TO MECHANICAL THROMBECTOMY?

¹JA Sousa, ¹R Machado, ¹L Cruz, ²J Paiva-Simões, ³L Santos-Martins, ¹L Almendra, ¹CC Nunes, ¹C Machado, ¹B Rodrigues, ¹O Galego, ¹C Nunes, ¹R Veiga, ¹E Machado, ¹F Silva, ¹G Santo, ¹⁻³J Sargento-Freitas. ¹Centro Hospitalar e Universitário de Coimbra, Coimbra; ²Centro Hospitalar e Universitário Lisboa Norte, Lisboa; ³Faculdade de Medicina da Universidade de Coimbra, Coimbra, Portugal

10.1136/neurintsurg-2021-ESMINT.40

Background In patients transferred from a primary stroke center (PSC) to a comprehensive stroke center (CSC) with a large

vessel occlusion (LVO-AIS) it is unclear who should undergo CT scan before mechanical thrombectomy (MT). We aimed to identify which patients benefited from such reevaluation.

Methods We conducted a retrospective study, including consecutive patients admitted to our CSC from 1/1/2016 to 31/12/2019, transferred from a PSC with anterior circulation LVO-AIS. We compared two groups. Group A (= beneficial CT scan) including patients either excluded from MT due to large infarction on CT reevaluation or patients headed directly to angio-suite and 90 day mRS >2. Group B (= futile CT scan) including patients submitted to MT and 90 day mRS ≤2 with or without prior CT reevaluation. We analyzed clinical and imaging variables collected in PSC. Collateral circulation was scored using modified Tan score.

Results We included 230 patients in group A and 225 in group B. Age (OR:1.04 95% CI 1.02–1.07, p<0.001), NIHSS (OR: 1.11 95%CI 1.06–1.15, p<0.001), ASPECTS score in PSC (OR:0.74 95%CI 0.60–0.91, p=0.006) and collateral score (OR:0.14 95%CI 0.083–0.222, p<0.001) were independent predictors of usefulness in CT scan reevaluation. Optimal cut-off points to identify best candidates for CT reevaluation were age ≥80, NIHSS ≥18 and ASPECTS ≤8.

Conclusion Patients with age ≥80, NIHSS ≥18, ASPECTS ≤8 or reduced collateral score (modified Tan <2) might benefit from CT scan reevaluation after transfer and prior to MT.

REFERENCES

1. Psychogios MN, et al. One-stop management of acute stroke patients minimizing door-to-reperfusion times. *Stroke* 2017.
2. Jadhav AP, et al. Interfacility transfer directly to the neuroangiography suite in acute ischemic stroke patients undergoing thrombectomy. *Stroke* 2017.
3. Mendez B, et al. Direct transfer to Angio-suite to reduce workflow times and increase favorable clinical outcome a case-control study. *Stroke* 2018.
4. Requena M, et al. Time matters: adjusted analysis of the influence of direct transfer to angiography-suite protocol in functional outcome. *Stroke* 2020.

Disclosure Nothing to disclose

EP42*

USE OF THE NEW PHILIPS' ADVANCED 3D NEURO-IMAGING FOR LVO TRIAGING AND DOOR TO PUNCTURE TIMES REDUCTION

¹F Massari, ¹J Singh, ¹AL Kuhn, ¹V Naragum, ¹MJ Gounis, ²B Hallacoglu, ¹A Puri. ¹University of Massachusetts Medical School, Worcester, MA, USA; ²Image Guided Therapy, Phillips Healthcare, Best, The Netherlands

10.1136/neurintsurg-2021-ESMINT.41

Introduction First clinical utilization of Philips' advanced 3D neuro-imaging prototype technology during stroke treatment.

Objectives and Aims Use of the new Philips' advanced 3D neuro-imaging prototype technology and cone beam angio-suite CTA versus standard multi-detector CT for triaging of acute LVO.

Methods We utilized the Philips' advanced 3D neuro-imaging prototype technology for within AngioLab non-contrast CT and CTA acquisition during direct to angio-suite LVO cases transfer. Pre-procedural ASPECT score and post-procedural evaluation of IPH or SAH components has been performed with the Philips' advanced 3D neuro-imaging prototype technology. The door-to-puncture timing in these patients has also been recorded and compared with that associated to standard patients necessitating of an additional stop in the ED for further/repetitive imaging acquisition.