



OPEN ACCESS

Original research

Cerebral venous outflow profiles are associated with the first pass effect in endovascular thrombectomy

Noel van Horn ,¹ Jeremy J Heit ,² Reza Kabiri,¹ Marius M Mader,³ Soren Christensen,⁴ Michael Mlynash,⁴ Gabriel Broocks ,¹ Lukas Meyer ,¹ Jawed Nawabi,⁵ Maarten G Lansberg,⁴ Gregory W Albers,⁴ Max Wintermark,² Jens Fiehler ,¹ Tobias D Faizy^{1,2}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/neurintsurg-2021-018078>).

¹Department of Diagnostic and Interventional Neuroradiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

²Department of Radiology, Stanford University School of Medicine, Stanford, California, USA

³Department of Neurosurgery, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

⁴Department of Neurology and Neurological Sciences, Stanford, Stanford, California, USA

⁵Department of Radiology, Charité School of Medicine and University Hospital Berlin, Berlin, Germany

Correspondence to

Dr Noel van Horn, Department of Diagnostic and Interventional Neuroradiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; no.vanhorn@uke.de

Received 30 July 2021

Accepted 24 October 2021

Published Online First

8 November 2021



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: van Horn N, Heit JJ, Kabiri R, *et al.* *J NeurolIntervent Surg* 2022;**14**:1056–1061.

ABSTRACT

Background Recent studies found that favorable venous outflow (VO) profiles are associated with higher reperfusion rates after mechanical thrombectomy (MT) in patients with acute ischemic stroke due to large vessel occlusion (AIS-LVO). Fewer retrieval attempts and first-pass revascularization during MT lead to better functional outcomes.

Objective To examine the hypothesis that favorable VO profiles assessed on baseline CT angiography (CTA) images correlate with successful vessel reperfusion after the first retrieval attempt and fewer retrieval attempts.

Methods A multicenter retrospective cohort study of patients with AIS-LVO treated by MT. Baseline CTA was used to determine the cortical vein opacification score (COVES). Favorable VO was defined as COVES ≥ 3 . Primary outcomes were successful with excellent vessel reperfusion status, defined as Thrombolysis in Cerebral Infarction (TICI) 2b/3 and 2c/3 after first retrieval attempt.

Results 617 patients were included in this study, of whom 205 (33.2%) had first pass reperfusion. In univariate analysis, ordinal COVES ($p=0.011$) values were significantly higher in patients with first pass than in those with non-first pass reperfusion, while the number of patients exhibiting favorable pial arterial collaterals using the Maas scale on CTA did not differ ($p=0.243$). In multivariable logistic regression analysis, higher COVES were independently associated with TICI 2b/3 (OR=1.25, 95% CI 1.1 to 1.42; $p=0.001$) and TICI 2c/3 (OR=1.2, 95% CI 1.04 to 1.36; $p=0.011$) reperfusion after one retrieval attempt, controlling for penumbra volume and time from symptom onset to vessel reperfusion.

Conclusions Favorable VO, classified as higher COVES, is independently associated with successful and excellent first pass reperfusion in patients with AIS-LVO treated by endovascular thrombectomy.

INTRODUCTION

The effectiveness of mechanical thrombectomy (MT) for the treatment of acute ischemic stroke due to large vessel occlusion (AIS-LVO) has been shown in several randomized trials.^{1–4} Successful flow restoration after the first retrieval attempt during MT (first pass effect, FPE) is associated with smaller final infarct volumes and better functional outcomes.⁵ In addition, recent studies have shown that an increased number

of retrieval attempts correlate with impaired functional outcomes after MT.⁶ Probably, these effects are not based on the FPE alone but may also be governed by robust collateral blood flow to and through the ischemic brain.^{7,8}

However, studies on the association of vascular factors that govern successful vessel reperfusion after the first retrieval attempt are scarce. Favorable arterial collateral blood flow to the ischemic tissue is probably correlated with successful vessel reperfusion after MT.^{9–10} However, recent studies reported that the assessment of venous outflow (VO) profiles may be more sensitive for the evaluation of collateral blood flow and tissue perfusion in patients with AIS-LVO expected to receive MT treatment, as it reflects blood flow after permeating the brain tissue.^{7,11,12} Favorable VO profiles were found to yield higher reperfusion rates after MT,¹³ but a correlation between VO profiles and the number of retrievals attempts during MT has not been investigated.

We hypothesized that favorable VO profiles in patients with AIS-LVO are associated with successful (Thrombolysis in Cerebral Infarction (TICI) score of 2b/3 and excellent (TICI 2c/3) vessel reperfusion after the FPE. We further hypothesized that patients with more favorable VO were likely to achieve reperfusion with fewer retrieval attempts during MT. We tested this hypothesis by determining VO using the cortical vein opacification score (COVES)⁷ on baseline CT angiography (CTA) images in patients with AIS-LVO treated by MT.

METHODS

Study population

In this multicenter retrospective cohort study, we analyzed 841 consecutive patients with AIS-LVO treated by MT at two comprehensive stroke centers (REDACTED) between October 2010 and September 2020. All patients with AIS-LVO routinely underwent MT according to standard of care. MT was withheld only in patients with a large early infarction (Alberta Stroke Program Early CT Score (ASPECTS) < 5) or a prolonged time window without evidence of perfusion mismatch.

Standard protocol approvals, registrations, and patient consent

All patient data obtained, and procedures conducted in this study, were obtained on approval by the institutional review boards of both stroke centers and complied with the Health Insurance Portability and Accountability Act (HIPAA) and followed the guidelines of the Declaration of Helsinki. Patient informed consent was waived by our review boards for this retrospective study.

Patient selection, population and clinical data

From each prospectively maintained stroke database, relevant clinical, imaging, and demographic data were obtained by reviewing electronic medical records. The MT procedure was performed using the local protocols and standard of care. If vessel reperfusion could not be achieved, rescue treatment included an additional combined aspiration/machinal thrombectomy maneuver with the option of a different stent retriever. The inclusion criteria were: (1) patients with AIS-LVO who underwent MT within 16 hours after symptom onset; (2) available baseline non-contrast head computed tomography and CTA with uniform opacification of the superior sagittal, transverse and sigmoid sinuses to the level of the jugular bulb to allow for VO determination; (3) evaluable CT perfusion (CTP) imaging with a core volume of <70 mL and a target mismatch ratio of >1.2; (4) large vessel occlusion of the anterior circulation affecting the internal carotid artery or first (M1) or second (M2) segment of the middle cerebral artery (MCA); (5) interpretable digital subtraction angiography images that allowed for a final TICI score assignment after MT; (6) documentation of the number of retrieval attempts during MT. Exclusion criteria were as follows: (1) non-evaluable CTA, due to poor image quality resulting from motion artifacts or insufficient opacification of the dural venous sinuses to allow for VO determination; (2) poor CTP image quality due to patient motion or failed contrast bolus; (3) absence of at least one follow-up head non-contrast CT/MRI examination within 72 hours of thrombectomy or hospital admission; (4) incomplete medical record data for TICI status and number of retrieval attempts performed during MT.

Image analysis

CTP images were automatically processed using RAPID (iScheMaView, Menlo Park, California, USA). The ischemic core was defined as the volume of tissue with at least a 70% reduction in cerebral blood flow relative to the contralateral cerebral hemisphere on CTP.

Venous outflow profiles were assessed by the COVES, quantifying the grade of opacification on a scale of 0 to 2 points (0=not visible; 1=moderate opacification; 2=full) for three major veins reflecting venous outflow of the MCA territory—namely, the vein of Labbé, sphenoparietal sinus, and superficial middle cerebral vein on single-phase baseline CTA.⁷ Favorable VO (VO+) profiles were defined as COVES of 3–6 and unfavorable VO (VO–) profiles were regarded as COVES of 0–2, based on previously published thresholds.¹² At first, VO classification was performed on a subset of image data by two neuroradiologists (JJH and TDF), one with 12 and one with 10 years of experience, to determine inter-rater agreement, followed by a rating of the remaining image data by one of these neuroradiologists (TDF). Intracranial pial arterial collaterals were evaluated by the Maas system.¹⁴ A

Table 1 Patient demographic, stroke presentation, and treatment details of 617 patients who underwent mechanical thrombectomy*

	First pass reperfusion (n=205)	Non-first pass reperfusion (n=282)	No reperfusion (n=130)	P value (FP vs non-FP)
Demographics				
Age (years), median (IQR)	77 (65–83)	74 (63–82)	77 (67–84)	0.183
Female	111 (54.1)	142 (50.4)	71 (54.6)	0.408
Medical history				
Atrial fibrillation	89 (43.6)	113 (40.5)	35 (27.1)	0.492
Hypertension	139 (68.1)	195 (69.6)	91 (70.5)	0.724
Hyperlipidemia	52 (25.4)	58 (20.6)	34 (26.2)	0.166
Diabetes mellitus	34 (16.7)	51 (18.2)	25 (19.4)	0.659
Blood glucose mg/dL, median (IQR)	120 (105–147)	119 (103–147)	123 (106–142)	0.720
Smoking history				
Current smoker	131 (64)	199 (70.6)	95 (73.1)	
Never smoked	26 (12.7)	30 (10.6)	13 (10.4)	
Prior smoker	29 (14.1)	36 (12.8)	17 (13.1)	
Stroke presentation details				
Presentation NIHSS score, median (IQR)	14 (8–18)	15 (9–19)	15 (10–19)	0.180
Onset to reperfusion (min), mean (SD)	110 (191)	133 (220)	120 (159)	0.003
Stroke treatment details				
IV tPA	106 (52.2)	139 (50)	66 (52)	0.631
Endovascular treatment details				
Aspiration	11 (5.4)	14 (5)	19 (14.6)	0.653
Aspiration and stent retriever	189 (92.2)	256 (90.7)	108 (83.1)	0.437
Balloon	5 (2.4)	9 (3.2)	3 (2.3)	0.361

Table 1 displays patients' demographic, stroke presentation and treatment findings of all 617 patients who underwent mechanical thrombectomy. Univariate logistic regression analysis between first pass- and non-first pass-reperused patients is displayed on the very right column with corresponding p values.
*Data are No. (N) and frequencies (%), unless otherwise indicated.
†Reperfusion is defined as Thrombolysis in Cerebral Infarction 2b/3.
FP, first pass reperfusion; NIHSS, National Institutes of Health Stroke Scale; Non-FP, non-first pass reperfusion; IV tPA, intravenous tissue plasminogen activator.

favorable collateral Maas score was defined either as arterial filling of the MCA territory equally to the contralateral hemisphere or greater than the contralateral normal side or when arterial collaterals were found to be exuberantly present.¹⁵

ASPECTS¹⁶ was independently determined on pretreatment non-contrast CT images by one neuroradiologist (TDF).

The angiographic results were assessed independently by two neuroradiologists, who were blinded to the ASPECTS readings, in the same manner as VO determination, on final digital subtraction angiography (DSA) image series for patients who underwent MT with or without previous intravenous administration of recombinant tissue plasminogen activator and was classified by the TICI scale; successful reperfusion was defined as TICI 2b/3 and excellent vessel reperfusion was regarded as TICI 2c/3. Any disagreement was settled by a consensus reading. The number of retrieval attempts was taken from the medical record data of the thrombectomy procedure. The specific thrombectomy techniques used in each group of patients are listed in [table 1](#).

Outcome measures

Primary outcomes were successful (TICI 2b/3) and excellent (TICI 2c-3) vessel reperfusion status after the first retrieval attempt on a final DSA series.

Statistical analysis

Statistical analyses were performed using SPSS Version 24.0 (IBM Inc., Armonk, New York, USA). Data are presented as mean±SD for normally distributed variables, as median and IQR for continuous variables, and as numbers and percentages for categorical variables. Non-reperfused (TICI 0 to 2a) patients were excluded for univariate regression analysis for primary outcome using χ^2 test for counts and Mann–Whitney U test for measurements. In univariable linear regression analysis proportional-odds assumption had to be met ($p>0.05$) before further analysis. Multivariable binary logistic regression analysis was performed to estimate the independent impact of COVES on first pass reperfusion at TICI 2b/3 and TICI 2c/3. Both models were adjusted for variables showing significance in the univariate analysis, that is, time from symptom onset to reperfusion, and penumbra volume ($T_{max} >6s$). Owing to a reportedly strong inter-relation of arterial clot localization and VO profiles,¹⁷ a variance of inflation analysis was conducted to investigate the covariance between arterial clot localizations and VO profiles in the respective regression models. Variance of inflation was found to be 7.34 and thus a high collinearity between VO and clot localization was assumed, which is in accordance with previous studies.¹⁷ Since the impact of clot localization on VO profiles has been investigated before and the primary focus of our analysis was to investigate the correlation between venous collateral biomarkers on FPE, we decided to exclude clot localization from the regression models under these circumstances, as it might introduce a high statistical bias to the interpretability of these models. Any unknown data were indicated at the given section below the tables.

RESULTS

Baseline characteristics of patients

A total of 617 patients met inclusion criteria (online supplemental figure 1) with a median age of 76 (IQR 64–83), 324 women (52.5%), a median National Institutes of Health Stroke Scale (NIHSS) score of 15 (IQR 9–19) and a median ASPECTS on admission of 8 (IQR 6–9). Of these patients, 205 (33.2%) experienced first pass reperfusion, while 282 (45.7%) needed at least one further retrieval attempt, and in 130 (21.1%) patients no vessel reperfusion was achieved during MT.

Interventional findings and treatment

Since our retrospective study spans a long period of time, detailed information about successful first pass reperfusions during the respective time is given in online supplemental table 1).

Comparison of patients with first pass and non-first pass reperfusion, showed no significant difference for age, admission NIHSS score, and admission ASPECTS ($p=0.183$, $p=0.18$, $p=0.42$, respectively). Similarly, frequency of intravenous tissue plasminogen activator administration was not different between both groups ($p=0.63$). On the contrary, patients with first pass reperfusion showed a shorter time interval between symptom onset to vessel reperfusion, with a mean of 110 min (SD 191, $p=0.003$) compared with non-first pass reperfusion (mean 133 min (SD 220)) as demonstrated in table 1. Moreover, compared with non-first pass reperfusion, first pass reperfusion had a significantly higher median COVES (median 3 (IQR

1–3) vs median 2 (IQR 1–3), $p=0.011$). A higher percentage of patients exhibited a favorable VO profile (47.4%) in the first pass-reperfusion group. Patients in that group also had a smaller penumbral volume (at $T_{max} >6s$) with a median of 115 mL (IQR 69–169, $p=0.05$), and first pass-reperfusion was significantly associated with the location of vessel occlusion ($p<0.001$). There were no significant differences in the frequency of the thrombectomy techniques used between the groups (table 1). The number of patients exhibiting a favorable Maas score profile was not significantly ($p=0.243$) different between the two groups (table 2).

Furthermore, compared with non-first pass reperfusion, first pass-reperfusion patients had less severe neurological deficits 24 hours after symptom onset with a median NIHSS score of 9 compared with 12 ($p=0.005$), and showed a significantly ($p=0.014$) higher degree of reperfusion with 59.5% versus 46.1% showing a TICI score of 3 (table 3).

Predictors of first pass reperfusion

In the univariate regression analysis, time to reperfusion, median COVES, penumbral volume (at $T_{max} >6s$), location of vessel occlusion, TICI scores, and 24-hour NIHSS score showed significant differences between the first pass reperfusion and the non-first pass reperfusion groups (tables 1 and 2).

For primary outcome prediction, 548 patients were included in the multivariable logistic regression model. Higher COVES (OR=1.25, 95% CI 1.1 to 1.42, $p=0.001$) were independently associated with TICI 2b/3 first pass reperfusion during MT, adjusting for the time from symptom onset to reperfusion (in min) and penumbra volumes ($T_{max} >6s$) (table 4 and figure 1).

In an additional second model, we included 518 patients to assess the correlation between COVES and excellent vessel reperfusion (TICI 2c/3) after a single retrieval. We found that COVES was independently associated with excellent vessel reperfusion (OR=1.2, 95% CI 1.04 to 1.36; $p=0.011$), after controlling for penumbra volume ($T_{max} >6s$) and time from symptom onset to reperfusion (in min) (table 5).

DISCUSSION

In this study of patients with AIS-LVO, we found that higher COVES (indicative of favorable VO profiles) were independently associated with higher rates of successful vessel reperfusion (TICI 2b/3) and excellent vessel reperfusion (TICI 2c/3) status during MT after the first retrieval attempt. Furthermore, compared with those with non-first pass reperfusion, these patients had a shorter time interval from symptom onset to vessel reperfusion, showed smaller penumbral volume at baseline on CTP, and experienced less severe early neurological dysfunctions 24 hours after stroke. Interestingly, the number of patients exhibiting a favorable pial arterial collateral profile on baseline CTA images did not differ between the first pass reperfusion and the non-first pass-reperfusion group. Thereby, our study underlines the important role of a sufficient cerebral venous collateral circulation in patients with AIS-LVO for early vessel reperfusion, which is associated with better functional outcomes.³ Our findings may add to a recent study that reported direct correlations of favorable VO profiles with good clinical outcomes after MT.¹²

Our findings demonstrate that higher COVES are independently associated with first pass reperfusion and fewer retrieval attempts, while pial arterial collaterals did not seem to affect first pass reperfusion during MT in our cohort. The substantial role of robust cerebral arterial collaterals on successful reperfusion during MT in patients with AIS-LVO has been demonstrated by prior studies.^{3,9} On the contrary, some findings suggest that pial arterial collaterals

Table 2 Imaging characteristics of 617 patients with a stroke*

	First pass reperfusion (n=205)	Non-first pass reperfusion (n=282)	No reperfusion (n=130)	P value (FP vs non-FP)
ASPECTS, median (IQR)	8 (7–9)	8 (6–9)	7 (6–9)	0.420
Favorable collateral Maas score	145 (71.1)	185 (66.1)	83 (63.8)	0.243
COVES				0.011
0	26 (12.7)	41 (14.6)	24 (18.5)	
1	36 (17.6)	80 (28.6)	40 (30.4)	
2	39 (19.1)	57 (20.4)	35 (26.9)	
3	59 (28.8)	53 (18.7)	19 (14.6)	
4	37 (18.1)	34 (12.1)	8 (6.2)	
5	8 (3.9)	15 (5.4)	3 (2.3)	
6	0 (0)	2 (0.7)	1 (0.8)	
Favorable VO profiles (COVES \geq 3)	97 (47.4)	107 (37.9)	35 (26.7)	0.005
Unfavorable VO profiles (COVES <2)	108 (52.6)	175 (62.1)	95 (73.3)	<0.001
Baseline estimated ischemic core volume (CBF <30%) (mL), median (IQR)	8.5 (0–30)	10 (0–31)	13 (0–41)	0.478
Penumbra Tmax >6s volume (mL), median (IQR)	115 (69–169)	128 (75–187)	114 (62–160)	0.050
Location of vessel occlusion				<0.001
Internal carotid artery	25 (12.3)	76 (27.1)	33 (25.4)	
MCA 1 segment occlusion	138 (67.6)	169 (60.4)	67 (51.5)	
MCA 2/distal segment occlusion	41 (20.1)	35 (12.5)	30 (23.1)	

*Data are No. and frequencies (%), unless otherwise indicated. ASPECTS, Alberta Stroke Program Early CT Score; CBF, cerebral blood flow; CO, cortical vein opacification score; FP, first pass reperfusion; MCA, middle cerebral artery; Tmax, Time-to-maximum of the residue function; VO, venous outflow.

do not promote successful vessel reperfusion and were not found to be associated with clinical outcomes after MT.¹⁸

In this regard, previous studies have shown that a VO biomarker, such as COVES may be more sensitive for tissue microperfusion assessment and collateral resilience, as it represents blood drainage after brain parenchyma circulation.^{7 19 20} This assumption has recently been supported by a retrospective cohort study, which found that favorable VO profiles were highly sensitive for robust tissue-level collaterals, automatically assessed on perfusion imaging.¹²

Our study showed that higher COVES were associated with a successful first pass reperfusion and fewer retrieval attempts to achieve reperfusion in patients with AIS-LVO during MT, which adds to a prior study which reported that robust VO profiles are associated with excellent vessel reperfusion (TICI 2c/3).¹³ Therefore, two pathophysiological assumptions may be postulated: First, as already reported by Goyal *et al*,³ and in accordance with our results, a shorter time interval from symptom onset vessel reperfusion and a consequently smaller infarct volume are associated with a smaller number of retrieval attempts. In our study, higher COVES were associated with smaller infarct volumes at baseline. Furthermore, higher COVES were correlated with a reduced number of retrieval attempts, which was found to be a key factor for successful vessel reperfusion during MT, as recently described by Flottmann *et al*.⁶ This again may prevent related complications and severe adverse events that directly affect the patient's functional outcome.⁶ Second, a well-developed cerebral tissue microperfusion is governed by a robust arterial collateralization and subsequent efficient venous drainage. In cases of a thrombotic large vessel occlusion, a suction is created on the distal site of the occluded vessel, which, in combination with the generally improved collateral microperfusion, may enable an easier and earlier MT. It is conceivable that favorable pial arterial collateralization and

subsequent VO distal to the thrombus may lead to an increased intravascular pressure, facilitating vessel clearance during thrombectomy. These inter-relations are in accordance with our hypothesis that a robust cerebral tissue collateral microcirculation yields first pass reperfusion.

However, our results need to be interpreted within certain reservations. VO robustness has been reported to be strongly dependent on the distinct localization of the arterial thrombus.¹⁷ Thus, in the event of a more proximal vessel occlusion (eg, internal carotid artery or proximal M1 occlusions including the perforating branches), VO is expectedly hampered, presumably owing to the reduced amount of blood perpetrating the affected brain tissue, and lower COVES were reported for this group of patients.¹⁷ We too found a strong collinearity between arterial clot localization and VO profiles (variance of inflation=7.34), with COVES becoming insignificant when adding arterial clot localization into the multi-variable regression models. These observations consequently allow for the presumption that the assumed beneficial effects of favorable VO profiles on FPE (more robust tissue microperfusion and elevated 'intravasal pressure' distal to the thrombus) that we speculated on above, become strongly mitigated in the event of a more proximal vessel occlusion, which is in accordance with the deliberations of prior studies.^{17 21}

More studies on the relationship between VO profiles and distinct clot localizations with respect to thrombectomy success are needed to validate our assumptions.

Our findings showed that patients with first pass reperfusion had significantly less early neurological dysfunctions and a stronger median NIHSS score improvement after 24 hours compared with patients with non-first pass reperfusion (tables 1–3). With respect to a recently reported threshold of an eight-point NIHSS score improvement,²² as a surrogate for long-term functional outcome, it would be of great interest to correlate VO profiles with early

Table 3 Thrombectomy procedural and clinical outcomes of 617 patients*

	First pass reperfusion (n=205)	Non-first-pass reperfusion (n=282)	No reperfusion (n=130)	P value (FP vs Non-FP)
Procedural variables				
General anesthesia				0.388
Conscious sedation	122 (74.4)	156 (69)	77 (68.1)	
General	31 (18.9)	56 (24.8)	23 (20.4)	
Conversion	11 (6.7)	14 (6.2)	13 (11.5)	
Procedure time, mean (SD)	10.6 (8.8)	12.9 (9.6)	15.4 (8.8)	0.069
Revascularization outcomes				
TICI 0	0 (0)	0 (0)	71 (54.6)	
TICI 1	0 (0)	0 (0)	14 (10.8)	
TICI 2a	0 (0)	0 (0)	45 (34.6)	
TICI 2b	76 (37.1)	138 (48.9)	0 (0)	
TICI 2c	7 (3.4)	14 (5)	0 (0)	
TICI 3	122 (59.5)	130 (46.1)	0 (0)	
Clinical Outcome				
NIHSS score at 24 hours, median (IQR)	9 (3–16)	12 (4–19)	15 (9–21)	0.005
NIHSS delta	-3 (-8–0)	-1 (-6–0)	0 (-4–3)	0.045
Modified Rankin Scale				0.413
Modified Rankin Scale ≤2 at 90 days	86 (43.2)	106 (39.1)	20 (15.9)	
Modified Rankin Scale 3–5 at 90 days	81 (40.7)	109 (40.2)	61 (48.4)	
Mortality at 90 days	32 (16.1)	56 (20.7)	45 (35.7)	
Univariate logistic regression analysis between first-pass- and non-firstpass-reperused patients is displayed on the very right column with corresponding p-values. Values are displayed as absolute numbers and frequencies, mean ± SD or median and interquartile range (IQR).				
*Data are No. and frequencies (%), unless otherwise indicated.				
FP, first pass reperfusion; NIHSS, National Institutes of Health Stroke Scale; Non-FP, non-first pass reperfusion; TICI, Thrombolysis in Cerebral Infarction.;				

neurological improvement in patients with AIS-LVO treated by MT. At this point, our study did not show a significant difference in NIHSS score improvement between the two groups analyzed. Thus, other postinterventional factors, such as severe adverse events, pneumonia, or myocardial infarction, may affect long-term clinical outcome and need further investigation. Nevertheless, approaching treatment decision-making by considering the collateral profile including VO and aiming at first pass reperfusion may contribute to a clinical neurological improvement of patients with

Table 4 Multivariable logistic regression analysis for primary outcome (successful reperfusion (TICI 2b/3) after first pass retrieval) of first pass and non-first pass-reperused patients with a TICI score of 2b/3 within the first retrieval attempt after mechanical thrombectomy.

Predictors	First pass reperfusion		
	OR	95% CI	P value
COVES	1.25	1.1 to 1.42	0.001
Time from onset to reperfusion	0.9994	0.998 to 1	0.204
Penumbra volume Tmax >6s	0.998	0.996 to 1	0.071
Observations n=548			
N=57 patients had unknown times from symptom onset to reperfusion, while in n=12 patients penumbra volume was unknown.			
COVES, cortical vein opacification score; TICI, Thrombolysis in Cerebral Infarction.			




Figure 1 Multivariable logistic regression plot to demonstrate probability of first pass reperfusion (y axis) based on COVES (x axis). COVES, cortical vein opacification score.

AIS-LVO. However, our study showed no significant difference in long-term functional outcomes (modified Rankin Scale score at 90 days) between non-first pass and first pass reperusers. Nevertheless, a recently conducted study showed that favorable VO profiles were independently associated with functional independence after MT at 90 days' follow-up.¹² On that note, results from the DEFUSE 3 group indicated that good arterial collaterals are associated with smaller ischemic core volumes, but not with good functional outcomes.²³ Thus, further investigations are warranted to understand better the impact of the cerebral collateral blood flow on tissue perfusion, MT success, and neurological outcomes.

Our study has some limitations inherent to its retrospective non-randomized design, which may introduce bias. Furthermore, this study was conducted by two primary stroke centers, who routinely perform MT not necessarily based on the same protocols, which may limit generalizability. For COVES rating, only single-phase CTA scans were used and venous opacification may be significantly affected by image selection, including acquisition timing and contrast injection rate.^{11,12} We acknowledge that use of single-phase CTA may limit the assessability of venous contrast opacification, as reported previously.²⁴ The high inter-relatedness of arterial clot localization and VO profiles may introduce bias to COVES assessment. It is yet unclear whether a timely resolved assessment of VO profiles (eg, using perfusion imaging or multiphase CTA)

Table 5 Multivariable logistic regression analysis for additional primary outcome (excellent reperfusion (TICI 2c/3) after first pass retrieval)

Predictors	First-Pass reperfusion		
	OR	95% CI	p-value
COVES	1.2	1.04 to 1.36	0.011
Time from onset to reperfusion	0.999	0.999 to 1.001	0.443
Penumbra volume Tmax >6s	0.998	0.995 to 0.999	0.038
Observations n=518			
Multivariable regression analysis in table 5 demonstrates calculated OR, 95% CI and p-value of all available (n=518) first pass- and non-first pass-reperused patients with a TICI score of 2 c/3 within the first retrieval attempt after mechanical thrombectomy.			
N=75 patients had unknown times from symptom onset to reperfusion and in 24 patients the exact penumbra volume was unknown.			
COVES, cortical vein opacification score; ;TICI, Thrombolysis in Cerebral Infarction.			

might mitigate this bias, as these methods allow more time for the cortical venous vessels to be filled with contrast agent. Anatomical variations of venous collaterals are quite common, introducing potential bias to COVES assessment.⁷ Finally, bias may be introduced by changing definitions of the target mismatch profiles used for patient selection in this retrospective study.

CONCLUSION

A favorable cerebral venous outflow profile classified by higher COVES and robust pial arterial collaterals is independently associated with first pass reperfusion in patients with AIS-LVO during MT. Our findings warrant further investigation on cortical venous pathophysiology in patients with AIS-LVO undergoing MT.

Twitter Noel van Horn @noelvanhorn, Jeremy J Heit @JeremyHeitMDPHD, Max Wintermark @stanfordNRAD and Jens Fiehler @Fie0815

Contributors NvH, JJH, and TDF conceptualized the study; NvH, MMM, and TDF performed data analyses and wrote the manuscript; NvH, RK, SC, MM, GB, LM, and MGL performed data preparation. NvH, JN, GWA, and TDF composed the cohort. JJH, GB, MW, JF and TDF supervised the clinical aspects of the study. JJH, GWA, JF, and MW supervised the work and wrote the manuscript. All authors critically reviewed the manuscript and approved the final version. TDF acts as the guarantor of the study.

Funding TDF was funded by the German Research Foundation (DFG) for his work as a postdoctoral research scholar at Stanford University, Department of Neuroradiology (Project Number: 411621970).

Competing interests JJH reports consulting for Medtronic and MicroVention and medical and scientific advisory board membership for iSchemaView. GWA reports equity and consulting for iSchemaView and consulting from Medtronic. JF reports membership of the editorial board of this journal. The other authors report no conflicts.

Patient consent for publication Not applicable.

Ethics approval All patient data were collected and analyzed in compliance with the respective institutional review boards (Administrative Panel on Human Subjects in Medical Research, Stanford University Institutional Review Board, Stanford, California, USA - ID: 37209; and the Ethics Committee of the Hamburg Chamber of Physicians, Hamburg, Germany - ID: NCT03356392). Any identifying information was removed in accordance with the Health Insurance Portability and Accountability Act (HIPAA) and the guidelines of the Declaration of Helsinki. Patient informed consent was waived by our review boards for this retrospective study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Noel van Horn <http://orcid.org/0000-0001-5764-1982>

Jeremy J Heit <http://orcid.org/0000-0003-1055-8000>
Gabriel Broocks <http://orcid.org/0000-0002-7575-9850>
Lukas Meyer <http://orcid.org/0000-0002-3776-638X>
Jens Fiehler <http://orcid.org/0000-0001-8533-7478>

REFERENCES

- Albers GW, Marks MP, Kemp S, *et al*. Thrombectomy for stroke at 6 to 16 hours with selection by perfusion imaging. *N Engl J Med* 2018;378:708–18.
- Berkhemer OA, Fransen PSS, Beumer D, *et al*. A randomized trial of intraarterial treatment for acute ischemic stroke. *N Engl J Med* 2015;372:11–20.
- Goyal M, Demchuk AM, Menon BK, *et al*. Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med* 2015;372:1019–30.
- Campbell BCV, Mitchell PJ, Kleinig TJ, *et al*. Endovascular therapy for ischemic stroke with perfusion-imaging selection. *N Engl J Med* 2015;372:1009–18.
- Zaidat OO, Castonguay AC, Linfante I, *et al*. First pass effect. *Stroke* 2018;49:660–6.
- Flottmann F, Brekenfeld C, Broocks G, *et al*. Good clinical outcome decreases with number of retrieval attempts in stroke thrombectomy: beyond the first-pass effect. *Stroke* 2021;52:482–90.
- Jansen IGH, van Vuuren AB, van Zwam WH, *et al*. Absence of cortical vein opacification is associated with lack of intra-arterial therapy benefit in stroke. *Radiology* 2018;286:643–50.
- Bhaskar S, Bivard A, Parsons M, *et al*. Delay of late-venous phase cortical vein filling in acute ischemic stroke patients: associations with collateral status. *J Cereb Blood Flow Metab* 2017;37:671–82.
- Bang OY, Saver JL, Kim SJ, *et al*. Collateral flow predicts response to endovascular therapy for acute ischemic stroke. *Stroke* 2011;42:693–9.
- Bang OY, Saver JL, Buck BH, *et al*. Impact of collateral flow on tissue fate in acute ischaemic stroke. *J Neurol Neurosurg Psychiatry* 2008;79:625–9.
- Hoffman H, Ziechmann R, Swarnkar A, *et al*. Cortical vein opacification for risk stratification in anterior circulation endovascular thrombectomy. *J Stroke Cerebrovasc Dis* 2019;28:1710–7.
- Faizy TD, Kabiri R, Christensen S, *et al*. Favorable venous outflow profiles correlate with favorable tissue-level collaterals and clinical outcome. *Stroke* 2021;52:1761–7.
- Faizy TD, Kabiri R, Christensen S, *et al*. Association of venous outflow profiles and successful vessel reperfusion after thrombectomy. *Neurology* 2021;96:10.1212/WNL.0000000000012106. [Epub ahead of print: 05 May 2021].
- Maas MB, Lev MH, Ay H, *et al*. Collateral vessels on CT angiography predict outcome in acute ischemic stroke. *Stroke* 2009;40:3001–5.
- Yeo LLL, Paliwal P, Teoh HL, *et al*. Assessment of intracranial collaterals on CT angiography in anterior circulation acute ischemic stroke. *AJNR Am J Neuroradiol* 2015;36:289–94.
- Barber PA, Demchuk AM, Zhang J, *et al*. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. *The Lancet* 2000;355:1670–4.
- Faizy TD, Kabiri R, Christensen S, *et al*. Distinct intra-arterial clot localization affects tissue-level collaterals and venous outflow profiles. *Eur J Neurol* 2021. doi:10.1111/ene.15079. [Epub ahead of print: 23 Aug 2021].
- de Havenon A, Mlynash M, Kim-Tenser MA, *et al*. Results from DEFUSE 3: good collaterals are associated with reduced Ischemic core growth but not neurologic outcome. *Stroke* 2019;50:632–8.
- Liebeskind DS. Imaging the collateralome: a stroke renaissance. *Curr Opin Neurol* 2015;28:1–3.
- Liebeskind D. The collateralome: a novel conceptual framework of systems biology in cerebrovascular disorders. *Brain Circulation* 2015;1:3–8.
- Behme D, Kowoll A, Weber W, *et al*. M1 is not M1 in ischemic stroke: the disability-free survival after mechanical thrombectomy differs significantly between proximal and distal occlusions of the middle cerebral artery M1 segment. *J Neurointerv Surg* 2015;7:559–63.
- Meyer L, Broocks G, Bechstein M, *et al*. Early clinical surrogates for outcome prediction after stroke thrombectomy in daily clinical practice. *J Neurol Neurosurg Psychiatry* 2020;91:1055–9.
- Ad H, Mlynash M, Kim-Tenser MA, *et al*. Results from DEFUSE 3. *Stroke* 2019;50:632–8.
- Jansen IGH, van Vuuren AB, van Zwam WH, *et al*. Absence of cortical vein opacification is associated with lack of intra-arterial therapy benefit in stroke. *Radiology* 2018;286