

While more than 90% of ICA aneurysms are microsurgically treated, 70% of ophthalmic aneurysms are treated with an endovascular approach.

In long-term follow-up, 90% of our patients have GOS 4 and 5, 2% have GOS 3 and 8% have GOS 1 or 2; that is not different in the endovascular or microsurgical group.

Conclusion Treatment of ICA aneurysms requires precise anatomical evaluation and a multi-disciplinary approach.

Both endovascular and microsurgical approaches are effective and have the same result.

Disclosure of Interest Nothing to disclose

P055/49 **COMPARISON OF NEUROFORM ATLAS STENT ASSISTED COILING AND COILING ALONE IN RUPTURED INTRACRANIAL ANEURYSMS: A PROPENSITY SCORE MATCHING ANALYSIS**

Ho Jun Yi*, Dong-Seong Shin, Bum-Tae Kim. *Soonchunhyang University Bucheon Hospital, Neurosurgery, Bucheon, South Korea*

10.1136/jnis-2023-ESMINT.88

Introduction Although Neuroform Atlas stent is commonly used in stent-assisted coiling (SAC) to treat ruptured intracranial aneurysms (RIA), its safety and efficacy remain controversial.

Aim of Study To assess the safety and efficacy of SAC using Neuroform Atlas for treating RIA compared with coiling alone by performing a propensity score matching analysis.

Methods RIA treated with coiling alone and SAC between January 2017 and May 2021 were retrospectively reviewed. Demographics, periprocedural complication rates, angiographic outcomes, and clinical outcomes of the SAC using Neuroform Atlas group and the coiling-alone group were analyzed with 1:1 propensity score matching.

Results A total of 375 aneurysms were enrolled, and 274 (63.1%) aneurysms were treated with coiling alone. In total, 101 (26.9%) aneurysms were treated with SAC, and Neuroform Atlas stent was used in 71 aneurysms. In propensity score matching, the SAC using Neuroform Atlas group showed higher incidence of complete occlusion (69.0% vs 56.3%, $P = .029$), lower rate of recanalization (11.3% vs 25.4%, $P = .011$), and lesser need for retreatment (7.0% vs 16.9%, $P = .016$) compared with the coiling-alone group. However, there were no significant differences in periprocedural complications such as intraprocedural thrombosis or postprocedural cerebral infarct between the 2 groups.

Conclusion The use of Neuroform Atlas is safe and effective for SAC in RIA with comparable procedure-related complication rates but better angiographic outcome in comparison with coiling alone.

Disclosure of Interest This study was supported by the National Research Foundation of Korea (NRF-2021R1G1A1094797) and the Soonchunhyang University Research Fund. The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

P056/52 **WIRELESS MICROCATETER PULL BACK TECHNIQUE (REVERSE PARKING TECHNIQUE) FOR MORE STABLE AND SAFE CEREBRAL COIL EMBOLIZATION**

Youngjin Jung*. *Neurosurgery, Daegu, South Korea*

10.1136/jnis-2023-ESMINT.89

Introduction Cerebral aneurysm coiling is a complex procedure that requires precise placement of the microcatheter into the aneurysm.

Aim of Study We examined the use of a new technique called the Wireless Microcatheter Pulled Back Technique (the Reverse Parking Technique), to improve the stability and safety of microcatheter placement.

Methods From July 1, 2022 to December 31, 2022, a total of 76 cases at a tertiary hospital underwent cerebral aneurysm coiling, with 51 cases utilizing the Reverse Parking Technique. The technique involves using a roadmap technique to guide the microcatheter over the microwire and into position within the parent vessel adjacent to the aneurysm, slightly passing the aneurysm. The microwire is then pulled into the microcatheter, and any slack or redundancy in the microcatheter is removed by gently pulling back on it. The microcatheter is slowly pulled back, allowing the tip to flip into the aneurysm.

Results Of the 51 cases that used the Reverse Parking Technique, 46 cases (90%) were successful, with no intraprocedural aneurysm perforation. In 5 cases, the technique failed, and the push technique was used instead. In one case, a thromboembolic complication occurred, but there was no neurological sequela. Two cases experienced unexpected premature microcatheter kicks back, but they were repositioned into the aneurysm relatively easily.

Conclusion The Reverse Parking Technique is a safe and reliable method for placing the microcatheter inside the aneurysm, particularly for small and ruptured aneurysms. It may help improve the success rate and reduce complications during cerebral aneurysm coiling procedures.

Disclosure of Interest Nothing to disclose

P057/53 **ENDOASCUAR TREATMENT OF THE MIDDLE CEREBRAL ARTERY HILLTOP ANEURYSM**

¹Sang Kyu Park*, ²Woo Sung Lee, ³Joon Ho Jung. ¹*Gangnam Severance Hospital, Neurosurgery, Seoul, South Korea;* ²*Ewha Womans University Seoul Hospital, Seoul, South Korea;* ³*Gangnam Severance Hospital, Seoul, South Korea*

10.1136/jnis-2023-ESMINT.90

Introduction 1 segment of middle cerebral artery (MCA) aneurysm is a relatively rare clinical condition. However, due to its complex geometry and deep location, microsurgical treatment is challenging.

Aim of Study We performed this study to define a specific form of M1 aneurysm that can be safely and effectively treated through EVT as M1 hilltop aneurysm, and to report our treatment results.

Methods Of 757 MCA aneurysm between December 2017 and October 2021, 54 M1 segment aneurysms were treated with EVT, and these aneurysms were designated M1 hilltop aneurysms. Clinical and radiographic data, including aneurysm characteristics, endovascular techniques, angiographic outcome, procedure-related complications and clinical outcomes at the time of the last follow-up, were collected and reviewed retrospectively.

Results Treatments were successful in all 54 cases; 21 cases were treated with coiling and 33 cases with stent-assist coiling (SAC). Of the 54 cases, 50 (92.6%) cases were identified as wide-neck aneurysms. The neck of aneurysm incorporating branch vessel was found in 49 (90.7%) cases. Immediate post-procedural angiogram showed favorable occlusion in 32 (59.3%), incomplete occlusion in 22 (40.7%). There were 4 (7.4%) procedure-related complications including thromboembolism and internal carotid artery dissection, but there were no cases of permanent neurological impairment. Recurrence was significantly related to aneurysmal neck (OR 3.9, 95% CI 1.2 to 12.9, $p=0.025$).

Conclusion EVT for M1 hilltop aneurysms appears to be safe and efficacious, with low mid-term recurrence rate. However, long-term and large cohort study will be needed.

Disclosure of Interest Nothing to disclose

P058/65

TWO-YEAR FOLLOW-UP OF DISTAL UNRUPTURED INTRACRANIAL ANEURYSMS TREATED WITH A SURFACE-MODIFIED FLOW DIVERTER UNDER PRASUGREL MONOTHERAPY

¹Luis Henrique Castro-Afonso*, ¹João Paulo Machado, ¹Guilherme Nakiri, ²Thiago Abud, ³Lucas Monsignore, ³Rafael Freitas, ⁴Ricardo Oliveira, ⁴Benedicto Colli, ¹Daniel Abud. ¹University of São Paulo, Interventional Neuroradiology, Ribeirão Preto; ²Hospital Israelita Albert Einstein, Interventional Neuroradiology, São Paulo; ³University of São Paulo, Ribeirão Preto; ⁴University of São Paulo, Division of Neurosurgery, Ribeirão Preto

10.1136/jnis-2023-ESMINT.91

Introduction A new generation of modified surface FDs and monotherapy using new antiplatelets may reduce both ischemic and hemorrhagic complications during the treatment of intracranial aneurysms. Long-term outcomes of distal IA treated with FD under antiplatelet monotherapy are unknown.

Aim of Study The aim of the present pilot study was to assess the two-year follow-up safety and efficacy of p48 MW HPC in the treatment of unruptured distal IA under the use of prasugrel monotherapy during 6 months following by aspirin until 24 months.

Methods This was a single-center, prospective, pivotal, open, single-arm study. The primary (safety) endpoint was the absence of any new neurological symptoms after treatment until the 24-month follow-up. The primary (efficacy) endpoint was the incidence of complete aneurysm occlusion 24 months after treatment. The secondary (efficacy) endpoints were incidence of aneurysm dome reduction $\geq 50\%$, and incidence of aneurysm dome reduction $< 50\%$, 24 months after treatment.

Results Twenty-one patients harboring 27 distal aneurysms of the anterior circulation were included. No patient had symptoms from treatment until 24-month follow-up. Complete aneurysm occlusion occurred in 20 (74%) of 27 aneurysms at the 24-month follow-up. Two aneurysms (7.4%) had a dome reduction $\geq 50\%$, 2 aneurysms (7.4%) had a dome reduction $< 50\%$, and 3 aneurysms (11.1%) remained unchanged.

Conclusion In this pilot trial, the treatment of distal, unruptured intracranial aneurysms with a stent under monotherapy with prasugrel appeared to be safe and effective.

Disclosure of Interest Nothing to disclose

P059/78

ENDOTHELIZATION OF SURFACE MODIFIED FLOW-DIVERTERS (FDS): DOES SURFACE MODIFICATION DELAY HEALING?

Christopher Zoppo, Mark Epshtein, Robert King, Matthew Gounis, Vania Anagnostakou*. Department of Radiology, New England Center for Stroke Research, UMass Chan Medical School, Worcester, USA

10.1136/jnis-2023-ESMINT.92

Introduction Flow diversion for aneurysm treatment has been widely used. Biomimetic surface modifications such as the phosphorylcholine polymer (Shield surface modification) aim to reduce thrombogenicity of these devices, and potentially the need for dual-antiplatelet therapy. How this modification affects endothelial growth remains unclear.

Aim of Study To investigate in-vivo the longitudinal healing of Pipeline Shield, Vantage Shield, and bare metal Pipeline Flex devices in vivo with the hypothesis that there would be no differences between modified and bare devices.

Methods The common carotid arteries of ten rabbits were implanted with the 3 FD-stents mentioned. All devices were imaged with high-frequency optical coherence tomography (HF-OCT) and angiography at implant, 5, 10, 15 and 30 days to evaluate placement and tissue growth. At 30 days, the devices were explanted, and endothelial growth was assessed with scanning electron microscopy (SEM) at 5 locations along their length using a semi-quantitative score from 0 (no tissue growth) to 5 (complete coverage). An automated method was used to measure the average tissue growth thickness (ATGT) along the device based on the HF-OCT images.

Results At 5 days, no statistical difference of ATGT between devices was shown, with the Vantage having a slightly higher ATGT. At the remaining timepoints, there was no difference, or trend in ATGT between device types. On SEM, all devices scored at least a 4 or 5 at every location, and no differences were found between median or mean scores.

Conclusion Endothelial coverage is not affected by Shield surface modification nor the Vantage stent design.

Disclosure of Interest CZ, ME, RK: Nothing to disclose

VA, MG :Medtronic

P060/83

OUTCOMES AND SAFETY OF FLOW DIVERTER STENTS FOR CAROTICO-OPHTHALMIC ANEURYSMS

¹Siôn Roberts*, ²Aiknaath Jain, ³Fathallah Islim, ³Hamed Nejadhamzeeigilani, ³Nayyar Saleem, ³Tony Goddard, ³Tufail Patankar. ¹Harrogate District Foundation Trust, Harrogate, UK; ²London North West University Healthcare Trust, HA1 3UJ, UK; ³Leeds Teaching Hospital Trust, Leeds, UK

10.1136/jnis-2023-ESMINT.93

Introduction Carotid-ophthalmic aneurysms represent 0.3–1% intracranial and 0.9–6.5% ICA aneurysms. Optimal management is currently contentious.

Aim of Study Reviewing outcome and safety of flow diverters stents (FDS) for Carotid-Ophthalmic aneurysms.

Methods Retrospective data between 2015 and 2021 was collected from CRIS/PPM/PACS databases for 64 carotid-