

SUPPLEMENTAL MATERIAL

Table 1. Comparison of the use of intraprocedural medications stratified by the presence of infarct (n = 156).

Medications, n (%)	Infarct (n = 97)	Non-infarct (n = 59)	p-value
Intra-arterial			.423
Tirofiban	2 (2)	2 (3)	
Tissue plasminogen activator	0 (0)	1 (2)	
Intravenous			.115
Tirofiban	3 (3)	0 (0)	

Table 2. Comparison of characteristics of the distal emboli between infarct and non-infarct groups (n = 222).

Variable	Infarct (n = 129)	Non-infarct (n = 93)	p-value
Vessel occluded, n (%)			.164
Anterior cerebral artery	3 (2)	4 (4)	
Posterior cerebral artery	3 (2)	1 (1)	
MCA M2 segment	19 (15)	8 (9)	
MCA distal M2 segment	4 (3)	2 (2)	
MCA M3 segment	42 (33)	21 (23)	
MCA distal M3 segment	22 (17)	28 (30)	
MCA M4 segment	36 (28)	29 (31)	
Brain parenchyma location, n (%)			.436
Medial	3 (2)	4 (4)	
Frontal	17 (13)	16 (17)	
Parietal	68 (53)	41 (44)	
Temporal	28 (22)	27 (29)	
Occipital	4 (3)	2 (2)	
Insula	9 (7)	3 (3)	

Abbreviations: MCA, middle cerebral artery.