

## SUPPLEMENTAL MATERIAL

**Table 1.** Comparison of the use of intraprocedural medications stratified by the presence of infarct (n = 156).

<b>Medications, n (%)</b>	<b>Infarct (n = 97)</b>	<b>Non-infarct (n = 59)</b>	<b>p-value</b>
Intra-arterial			.423
Tirofiban	2 (2)	2 (3)	
Tissue plasminogen activator	0 (0)	1 (2)	
Intravenous			.115
Tirofiban	3 (3)	0 (0)	

**Table 2.** Comparison of characteristics of the distal emboli between infarct and non-infarct groups (n = 222).

Variable	Infarct (n = 129)	Non-infarct (n = 93)	p-value
<b>Vessel occluded, n (%)</b>			.164
Anterior cerebral artery	3 (2)	4 (4)	
Posterior cerebral artery	3 (2)	1 (1)	
MCA M2 segment	19 (15)	8 (9)	
MCA distal M2 segment	4 (3)	2 (2)	
MCA M3 segment	42 (33)	21 (23)	
MCA distal M3 segment	22 (17)	28 (30)	
MCA M4 segment	36 (28)	29 (31)	
<b>Brain parenchyma location, n (%)</b>			.436
Medial	3 (2)	4 (4)	
Frontal	17 (13)	16 (17)	
Parietal	68 (53)	41 (44)	
Temporal	28 (22)	27 (29)	
Occipital	4 (3)	2 (2)	
Insula	9 (7)	3 (3)	

Abbreviations: MCA, middle cerebral artery.